

The existential theory of equations with
rational constraints in free groups is
PSPACE-complete

Volker Diekert¹, Claudio Gutiérrez², Christian Hagenah¹

April 12, 2005

¹ Institut für Formale Methoden der Informatik (FMI),
Universität Stuttgart,
Universitätsstr. 38
D-70569 Stuttgart, Germany
diekert@fmi.uni-stuttgart.de christian@hagenah.de

² Depto. de Ciencias de la Computación, Universidad de Chile,
Blanco Encalada 2120, Santiago, Chile
cgutierr@dcc.uchile.cl

ACM Classification F.2. Analysis of Algorithms and Problem Complexity, F.2.2. Computation on Discrete Structures, F.4. Mathematical Logic and Formal Languages.

Subject Descriptor Equations in Free Groups.

Contents

1	Introduction	3
2	Basic Notions and Statements of Theorems	4
2.1	Preliminaries	4
2.2	Free Groups	5
2.3	Free Monoids with Involution	6
2.4	Equations with Constraints	7
3	Reductions	9
3.1	Reduction of Problem EFG to EFMI	9
3.2	Reduction of Problem EFMI to EWC	10
4	Problem EWC is in Pspace	12
4.1	Road-Map	12
4.2	A PSPACE-Complete Subproblem	14
4.3	The Exponent of Periodicity	15
4.4	Exponential Expressions	15
4.5	Base Changes	17
4.6	Projections	18
4.7	Partial Solutions	20
4.8	The Search Graph and Plandowski's Algorithm	22
4.9	Free Intervals	24
4.10	Critical Words	29
4.11	The ℓ -Transformation	31
4.12	The ℓ -transformation E_ℓ is Admissible	33
4.13	The Arc from E_0 to E_1	36
4.14	The Equation $E_{\ell,\ell'}$ for $1 \leq \ell < \ell' \leq 2\ell$	36
4.15	Passing from E_ℓ to $E_{\ell,\ell'}$ for $1 \leq \ell < \ell' \leq 2\ell$	39
5	Concluding Remarks	41
6	Appendix: Proof of Proposition ??	42
	Acknowledgments	45
	Bibliography	46

Abstract

It is well-known that the existential theory of equations in free groups is decidable. This is a celebrated result of Makanin which was published 1983. Makanin did not discuss complexity issues, but later it was shown that the scheme of his algorithm is not primitive recursive. In this paper we present an algorithm that works in polynomial space. This improvement is based upon an extension of Plandowski's techniques for solving word equations. We present a PSPACE-algorithm in a more general setting where each variable has a rational constraint, that is, the solution has to respect a specification given by a regular word language. We obtain our main result about the existential theory in free groups as a corollary of the corresponding statement in free monoids with involution.

1 Introduction

Around 1980 great progress was achieved on the algorithmic decidability of elementary theories of free monoids and groups. In 1977 Makanin [24] proved that the existential theory of equations in free monoids is decidable by presenting an algorithm which solves the satisfiability problem for a single word equation with constants. In 1983 he extended his result to the more complicated situation in free groups [25]. Using a result by Merzlyakov [31] Makanin also showed that the positive theory of equations in free groups is decidable [26], and Razborov was able to give a description of the whole solution set [37]. The algorithms of Makanin are very complex: For word equations the running time was first estimated by several towers of exponentials and it took more than 20 years to lower it to the best known bound for Makanin's original algorithm, which is EXPSPACE [14]. For solving equations in free groups Kościelski and Pacholski [21] have shown that the scheme proposed by Makanin is not primitive recursive.

In 1999 Plandowski used another method for solving word equations and he showed that the satisfiability problem for word equations is in PSPACE, [34, 35]. One ingredient of his work is to use data compression to reduce the space. The importance of data compression was first recognized by Rytter and Plandowski when applying Lempel-Ziv encodings to the minimal solution of a word equation [36]. Another important definition is the ℓ -factorization of a solution to a word equation. The roots of the notion of ℓ -factorization are in the notion of synchronizing factorization from [19]. Gutiérrez extended Plandowski's method to the case of free groups, [16]. Thus, a non-primitive recursive scheme for solving equations in free groups has been replaced by

a polynomial space bounded algorithm. Hagenah and Diekert worked independently in the same direction and using some ideas of Gutiérrez [15] they obtained a result which includes the presence of rational constraints [4, 17]. The present paper is a journal version of [4, 16]. It shows that the existential theory of equations in free groups with rational constraints is PSPACE-complete. Rational constraints mean that a possible solution has to respect a specification which is given by a regular word language. The idea to consider regular constraints for word equations goes back to Schulz [38] who also pointed out the importance of this concept, see also [7, 13]. The PSPACE-completeness for the case of word equations with regular constraints has already been stated by Rytter according to [34, Thm. 1].

Our proof reduces the case of equations with rational constraints in free groups to the case of equations with regular constraints in free monoids with involution, which turn out to be central objects. (Makanin uses the notion of “paired alphabet;” the main difference is that he considered “non contractible” solutions only, whereas we deal with general solutions. Allowing general solutions simplifies the presentation considerably.) Our work extends the method of [34, 35] so that it copes with involutions, and it extends the method of [16] so that it copes with rational constraints. The first step is a reduction to the satisfiability problem of a single equation with regular constraints in a free monoid with involution. In order to avoid an exponential blow-up, we do not use a reduction as in [26], but a simpler one. In particular, we can handle negations simply by positive rational constraints. In the second step we show that the satisfiability problem of a single equation with regular constraints in a free monoid with involution is in PSPACE. This part is technical and first we introduce several notions like base-change, projection, partial solution, and free interval. After these preparations we can follow Plandowski’s method. Throughout we shall use many of the deep ideas which were presented in [34, 35], but we apply them in a different setting. Hence, as we cannot use Plandowski’s result as a black box, we go through the whole construction again and we obtain a self-contained presentation.

2 Basic Notions and Statements of Theorems

2.1 Preliminaries

An *involution* on a set is a bijection $\bar{}$ such that $\overline{\overline{x}} = x$ for all elements x . If M is a monoid, then an involution $\bar{} : M \rightarrow M$ means that we also require $\overline{1} = 1$ for the unit element 1 and $\overline{xy} = \overline{y}\overline{x}$ for all $x, y \in M$.

Let Σ be a finite alphabet. By Σ^* we denote the free monoid over Σ . Elements

of Σ^* are called *words*. The length of a word w is denoted by $|w|$. A *factor* of a word w is a word v such that $w = w_1vw_2$. By $F(\Sigma)$ we denote the free group over Σ . Elements of $F(\Sigma)$ can be represented by words over $\Gamma = \Sigma \cup \bar{\Sigma}$, where $\bar{\Sigma} = \{\bar{a} \mid a \in \Sigma\}$ is a disjoint copy of Σ . We let $\bar{a} = a$, this defines an involution $\bar{\cdot} : \Gamma \rightarrow \Gamma$; and the involution is extended to Γ^* by $\overline{a_1 \cdots a_n} = \bar{a}_n \cdots \bar{a}_1$. The meaning of \bar{w} is the inverse w^{-1} in $F(\Sigma)$. A word $w \in \Gamma^*$ is *freely reduced*, if it contains no factor of the form $a\bar{a}$ with $a \in \Gamma$. For $w \in \Gamma^*$ we denote by \hat{w} the freely reduced word which denotes the same group element in $F(\Sigma)$. Hence, $\hat{u} = \hat{v}$ if and only if $\psi(u) = \psi(v)$, where $\psi : \Gamma^* \rightarrow F(\Sigma)$ denotes the canonical homomorphism.

The classes of *rational* and *recognizable* subsets are defined for every monoid M , [10]. Rational sets (or languages) are defined inductively as follows. All finite subsets of M are rational. If $C_1, C_2 \subseteq M$ are rational, then the union $C_1 \cup C_2$, the concatenation $C_1 \cdot C_2$, and the generated submonoid C_1^* are rational. A subset $C \subseteq M$ is recognizable, if and only if there is a homomorphism h to some finite monoid M' such that $C = h^{-1}h(C)$. Kleene's Theorem states that in finitely generated free monoids both classes coincide, and we follow the usual convention to call a rational (or recognizable) subset of a free monoid *regular*.

The empty word is the unit element of a free monoid, it is denoted by 1 as the unit element in other monoids. The singleton set $\{1\}$ is rational in $F(\Sigma)$, but not recognizable if $\Sigma \neq \emptyset$. A subset $C \subseteq F(\Sigma)$ is rational if and only if $C = \psi(C')$ for some regular language $C' \subseteq \Gamma^*$. In particular, we can use a non-deterministic finite automata over Γ for specifying rational group languages over $F(\Sigma)$.

The *existential theory of equations with rational constraints* in a monoid M with a generating set Γ is defined as follows. Let Ω be a set of variables (or unknowns). Atomic formulae are either of the form $L = R$, where $L, R \in (\Gamma \cup \Omega)^*$ or of the form $X \in C$, where X is in Ω and $C \subseteq M$ is a rational language. An *existentially quantified* formula is a block of existentially quantified variables followed by a Boolean combination of atomic formulae. It is closed, if there are no free variables. The existential theory of equations with rational constraints in M is the set of all closed existentially quantified formulae which are *true* over M .

2.2 Free Groups

The following proposition is due to Benois [1], see also [2, Sect. III.2].

Proposition 1 *The family of rational languages over the free group $F(\Sigma)$ forms an effective Boolean algebra.*

Proof. (Sketch.) It is enough to show that the family of rational languages is closed under complementation. Let $C' \subseteq \Gamma^*$ be a regular language and $C = \psi(C')$ the corresponding rational group language in $F(\Sigma)$. Assume that C' is given by some non-deterministic finite automaton. Using the same state set we can construct (in polynomial time) a finite automaton which accepts the following language

$$C'' = \{ v \in \Gamma^* \mid \exists u \in C' : u \xrightarrow{*} v \}$$

where $u \xrightarrow{*} v$ means that v is a descendant of u by the rewriting system $\{ a\bar{a} \rightarrow 1 \mid a \in \Gamma \}$. Then we complement C'' with respect to Γ^* ; and intersect $\Gamma^* \setminus C''$ with the regular set of freely reduced words. We obtain a regular set \tilde{C}' . Hence, the complement of C in $F(\Sigma)$ is the rational group language $\psi(\tilde{C}')$. \square

Problem 2 *By EFG we denote the following decision problem:*

INPUT: A finite alphabet Σ and a closed existentially quantified formula with rational constraints in the free group $F(\Sigma)$.

QUESTION: Is the formula true in $F(\Sigma)$?

Theorem 3 *The problem EFG is PSPACE-complete.*

The difficult part is to show that EFG is in PSPACE. For this we prove a more general statement about the existential theory of equations with regular constraints in free monoids with involution.

2.3 Free Monoids with Involution

In the following let Γ be a finite alphabet of constants and Ω be an alphabet of variables together with involutions $\bar{\cdot} : \Gamma \rightarrow \Gamma$ and $\bar{\cdot} : \Omega \rightarrow \Omega$. The involution on Ω is without fixed points, but we allow fixed points for the involution on Γ . The involution is extended to $(\Gamma \cup \Omega)^*$ by $\overline{x_1 \cdots x_n} = \bar{x}_n \cdots \bar{x}_1$ for $n \geq 0$ and $x_i \in \Gamma \cup \Omega$, $1 \leq i \leq n$. Clearly, $\overline{\bar{u}} = u$ for all $u \in (\Gamma \cup \Omega)^*$.

From now on, almost all monoids M under consideration are equipped with an involution $\bar{\cdot} : M \rightarrow M$. A *morphism* between monoids with involution M and M' is henceforth a mapping $h : M \rightarrow M'$ such that $h(1) = 1$, $h(xy) = h(x)h(y)$, and $h(\bar{x}) = \overline{h(x)}$ for all $x, y \in M$. Thus, a morphism is a homomorphism of monoids which respects the involution. The pair $(\Gamma^*, \bar{\cdot})$ is called a *free monoid with involution*. A morphism $h : \Gamma^* \rightarrow M$ is specified by a list $(h(a); a \in \Gamma)$ such that $h(\bar{a}) = \overline{h(a)}$ for all $a \in \Gamma$.

Problem 4 By EFMI we denote the following decision problem:

INPUT: A closed existentially quantified formula with regular constraints in a free monoid with involution $(\Gamma^*, \bar{})$.

QUESTION: Is the formula true in $(\Gamma^*, \bar{})$?

The proof of the following statement is the main technical contribution of the paper.

Theorem 5 The problem EFMI is PSPACE-complete.

2.4 Equations with Constraints

In the following it is more suitable to work with Boolean matrices instead of finite automata. Let $n \geq 1$. Henceforth, $M_{2n} \subseteq \mathbb{B}^{2n \times 2n}$ denotes the following monoid with involution:

$$M_{2n} = \left\{ \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \mid A, B \in \mathbb{B}^{n \times n} \right\},$$

where

$$\overline{\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}} = \begin{pmatrix} B & 0 \\ 0 & A \end{pmatrix}^T = \begin{pmatrix} B^T & 0 \\ 0 & A^T \end{pmatrix}$$

The operator T denotes transposition and $\mathbb{B}^{n \times n}$ is the monoid of Boolean $n \times n$ -matrices.

Definition 6 An equation E with constraints is a list

$$E = (\Gamma, h, \Omega, \rho; L = R)$$

containing the following items:

- The alphabet $(\Gamma, \bar{})$ with involution.
- The morphism $h : \Gamma^* \rightarrow M_{2n}$ which is specified by a mapping $h : \Gamma \rightarrow M_{2n}$ such that $h(\bar{a}) = \overline{h(a)}$ for all $a \in \Gamma$.
- The alphabet $(\Omega, \bar{})$ with involution without fixed points.
- A mapping $\rho : \Omega \rightarrow M_{2n}$ such that $\rho(\bar{X}) = \overline{\rho(X)}$ for all $X \in \Omega$.
- The word equation $L = R$ where $L, R \in (\Gamma \cup \Omega)^+$.

A solution of E is a mapping $\sigma : \Omega \rightarrow \Gamma^*$, which is extended to a morphism $\sigma : (\Gamma \cup \Omega)^* \rightarrow \Gamma^*$ by leaving the letters from Γ invariant such that the following three conditions are satisfied:

$$\begin{aligned} \sigma(L) &= \sigma(R), \\ \sigma(\overline{X}) &= \overline{\sigma(X)} \quad \text{for all } X \in \Omega, \\ h\sigma(X) &= \rho(X) \quad \text{for all } X \in \Omega. \end{aligned}$$

Let $d = |LR|$ be the denotational length of the word equation $L = R$. The input size of E is given by

$$\|E\| = d + n + \log_2(|\Gamma| + |\Omega|).$$

The definition of input size takes into account that there might be constants or variables with constraints which are not present in the equation. Due to this definition we assume that the input to Problem 7 is kept on a separate read-only storage.

Problem 7 By EWC we denote the following decision problem:

INPUT: An equation with constraints, $E = (\Gamma, h, \Omega, \rho; L = R)$.

QUESTION: Is there a solution $\sigma : \Omega \rightarrow \Gamma^*$?

Theorem 8 The problem EWC is PSPACE-complete.

We now turn to the proofs of Theorems 3, 5, and 8. The PSPACE-hardness of the problems EFMI, EFG, and EWC follows directly from a result of Kozen [22], since the *empty intersection* problem of regular sets can easily be encoded in the problems above. Therefore the PSPACE-hardness is not discussed further in the sequel.

The difficult part is to show that the problems EFG, EFMI, and EWC can be solved in polynomial space. We proceed as follows. Section 3.1 yields a (polynomial time) reduction from the problem EFG to the problem EFMI. Section 3.2 yields a reduction from EFMI to EWC, but this reduction involves non-deterministic steps. It can be performed however in non-deterministic polynomial time. Section 4 is the core of the paper. It shows that the problem EWC can be solved by some non-deterministic PSPACE algorithm. By Savitch's theorem such a procedure can be transformed into a polynomially space bounded deterministic decision procedure, see e.g. [18]. This concludes the proof of Theorems 3, 5, and 8.

Remark 9 Problem EWC is NP-hard for $n = 1$ already, since then we are in the framework of word equations (without constraints); and linear integer programming can easily be reduced to word equations, see e.g. [3]. We conjecture that the problem is in fact NP-complete, if n is bounded by some constant which is not part of the input, see also [36].

3 Reductions

3.1 Reduction of Problem EFG to EFMI

The next technical lemma follows directly from the well known fact that the Cayley graph of a free group is a tree. The proof of Lemma 10 is therefore omitted. As above, let $\psi : \Gamma^* \rightarrow F(\Sigma)$ be the canonical morphism.

Lemma 10 *Let $u, v, w \in \Gamma^*$ be freely reduced words. Then we have $uvw = 1$ in $F(\Sigma)$ (i.e. $\psi(uvw) = 1$) if and only if there are words $P, Q, R \in \Gamma^*$ such that $u = P\bar{Q}$, $v = Q\bar{R}$, and $w = R\bar{P}$ in Γ^* .*

Proposition 11 *There is a polynomial time reduction of problem EFG to EFMI.*

Proof. The reduction follows standard lines. The input to the problem EFG is a closed existentially quantified formula with rational constraints in the free group $F(\Sigma)$. Using De Morgan's law we may assume that there are no negations at all. Since we are in a group, the atomic formulae are now of the either form: $W = 1$, $W \neq 1$, $X \in C$ or $X \notin C$ where $W \in (\Gamma \cup \Omega)^*$, $X \in \Omega$, and $C \subseteq F(\Sigma)$ is rational. The reason that we keep $X \notin C$ instead of $X \in \tilde{C}$ where $\tilde{C} = F(\Sigma) \setminus C$ is that the complementation may involve an exponential blow-up.

The next step is to replace every formula $W \neq 1$ by

$$\exists X : WX = 1 \wedge X \notin \{1\},$$

where X is a fresh variable, hence we can put $\exists X$ to the front.

We may assume that $|W| \geq 3$, since if $1 \leq |W| < 3$, then we may replace $W = 1$ by $W a \bar{a} = 1$ for some $a \in \Gamma$. For the present reduction it is convenient to assume that $|W| = 3$ for all subformulae $W = 1$. This is easy to achieve. As long as there is a subformula $x_1 \cdots x_k = 1$, $x_i \in \Gamma \cup \Omega$ for $1 \leq i \leq k$ and $k \geq 4$, we replace it by the conjunction

$$\exists Y : x_1 x_2 Y = 1 \wedge \bar{Y} x_3 \cdots x_k = 1,$$

where Y is a fresh variable and $\exists Y$ is put to the front, and then proceed recursively.

Now, there are no negations and all atomic formulae are of type $W = 1$, $X \in C$ or $X \notin C$, where $W \in (\Gamma \cup \Omega)^+$, $|W| = 3$, $X \in \Omega$, and $C \subseteq F(\Sigma)$ is rational.

At this point we switch to free monoids with involution. Recall that $\psi : \Gamma^* \rightarrow F(\Sigma)$ denotes the canonical morphism and that $X \in C$ (resp. $X \notin C$) means

in fact $X \in \psi(C')$ (resp. $X \notin \psi(C')$), where $C' \subseteq \Gamma^*$ is a regular language specified by some finite non-deterministic automaton over the alphabet Γ^* . Using ψ -symbols we obtain an interpretation over (Γ^*, \neg) without changing the truth value of the input formula: We replace each subformula $X \in C$ (resp. $X \notin C$) syntactically by $\psi(X) \in \psi(C')$ (resp. $\psi(X) \notin \psi(C')$) and we replace each subformula $W = 1$ by $\psi(W) = 1$.

We keep the interpretation over words, but we now eliminate all occurrences of ψ again. We begin with the occurrences of ψ in the constraints. Let $C' \subseteq \Gamma^*$ be regular. According to the proof of Proposition 1 we construct a finite automaton, which accepts the following language

$$C'' = \{ v \in \Gamma^* \mid \exists u \in C' : u \xrightarrow{*} v \}.$$

In particular, $\psi(C') = \psi(C'')$ and $\widehat{C} \subseteq C''$ where $\widehat{C} = \{ \widehat{u} \in \widehat{\Gamma^*} \mid u \in C' \}$. We replace all positive atomic subformulae of the form $\psi(X) \in \psi(C')$ by $X \in C''$. A simple reflection shows that the truth value has not changed since we can think of X as being a freely reduced word. For a negative formula $\psi(X) \notin \psi(C')$ we have to be a little more careful. Let $N \subseteq \Gamma^*$ be the regular set of all freely reduced words. The language N is accepted by some deterministic finite automaton with $|\Gamma| + 2$ states. We replace $\psi(X) \notin \psi(C')$ by

$$X \notin C'' \wedge X \in N,$$

where C'' is as above. Again the truth value did not change.

We now have to deal with the formulae $\psi(xyz) = 1$ where $x, y, z \in \Gamma \cup \Omega$. Observe that the underlying quantifier free formula is satisfiable over Γ^* if and only if it is satisfiable in freely reduced words.

Based on Lemma 10 we replace each atomic subformulae $\psi(xyz) = 1$ with $x, y, z \in \Gamma \cup \Omega$ by a conjunction

$$\exists P \exists Q \exists R : x = P\overline{Q} \wedge y = Q\overline{R} \wedge z = R\overline{P},$$

where P, Q, R are fresh variables and the existential block is put to the front. The new existential formula has no occurrence of ψ anymore. The atomic subformulae are of the form $x = yz$, $X \in C$ or $X \notin C$, where $x, y, z \in \Gamma \cup \Omega$ and $C \subseteq \Gamma^*$ is regular. The size of the new formula is polynomial in the size of the original formula. This finishes the reduction from the problem EFG to EFMI. \square

3.2 Reduction of Problem EFMI to EWC

Proposition 12 *There is a non-deterministic polynomial time reduction of problem EFMI to EWC.*

Proof. The input to problem EFMI is a closed existentially quantified formula Φ with regular constraints over a free monoid with involution. We define a procedure which transforms the input Φ into an equation with constraints E_Φ . If Φ is true, then at least one possible output E_Φ has a solution. If the output E_Φ has a solution, then Φ is true. The procedure will work in non-deterministic polynomial time.

We may assume that the formula Φ contains no negations and all atomic subformulae are of type $U = V$, $U \neq V$, $X \in C$ or $X \notin C$, where $U, V \in (\Gamma \cup \Omega)^*$, $X \in \Omega$, and $C \subseteq \Gamma^*$ is regular.

Since we work over a free monoid Γ^* it is easy to handle inequalities $U \neq V$ where $U, V \in (\Gamma \cup \Omega)^*$. If two words u, v in Γ^* are different, then there are three cases: u is a proper prefix of v or v is a proper prefix of u or there is some word x such that xa is a prefix of u , xb is a prefix of v , and $a \neq b$. Therefore, a subformula $U \neq V$ can be replaced by

$$\exists X \exists Y \exists Z : \bigvee_{a \in \Gamma} (U = VaX \vee V = UaX \vee \bigvee_{a \neq b \in \Gamma} (U = XaY \wedge V = XbZ)).$$

Making guesses we can eliminate all disjunctions to obtain an existentially quantified formula which consists of a block of existentially quantified variables followed by a single conjunction over atomic subformulae of type $U = V$, $X \in C$ or $X \notin C$, where $U, V \in (\Gamma \cup \Omega)^*$, $X \in \Omega$, and $C \subseteq \Gamma^*$ is regular. By a standard procedure we can replace a conjunction of word equations over $(\Gamma \cup \Omega)^*$ by a single word equation $L = R$ where neither L nor R is empty. For example, we may choose a new letter a ($a \notin \Gamma$) and then we can replace a system $L_1 = R_1, L_2 = R_2, \dots, L_k = R_k$ by $L_1 a L_2 a \dots a L_k = R_1 a R_2 a \dots a R_k$ and we add for all variables X the constraint $X \in \Gamma^*$.

Therefore, we may assume that our input is now given by three items: a single word equation $L = R$ with $L, R \in (\Gamma \cup \Omega)^+$ and two lists: $(X_j \in C_j, 1 \leq j \leq m)$ and $(X_j \notin C_j, m < j \leq k)$. Each regular language $C_j \subseteq \Gamma^*$ is specified by some non-deterministic automaton $\mathcal{A}_j = (Q_j, \Gamma, \delta_j, I_j, F_j)$ where Q_j is the set of states, $\delta_j \subseteq Q_j \times \Gamma \times Q_j$ is the transition relation, $I_j \subseteq Q_j$ is the subset of initial states, and $F_j \subseteq Q_j$ is the subset of final states, $1 \leq j \leq k$. Of course, a variable X may occur several times in the list with different constraints, therefore we might have k greater than $|\Omega|$.

For the reduction to the problem EWC we have to consider Boolean matrices instead of finite automata. This allows us to store all constraints concerning a variable in a single Boolean matrix. Let Q be the disjoint union of the state spaces Q_j , $1 \leq j \leq k$. We may assume that $Q = \{1, \dots, n\}$. Let $\delta = \bigcup_{1 \leq j \leq k} \delta_j$, then $\delta \subseteq Q \times \Gamma \times Q$ and with each $a \in \Gamma$ we can associate a Boolean $n \times n$ matrix $g(a) \in \mathbb{B}^{n \times n}$ such that $g(a)_{i,j} = 1$, if $(i, a, j) \in \delta$ and

$g(a)_{i,j} = 0$ otherwise. We define a morphism $h : \Gamma^* \rightarrow M_{2n}$ by

$$h(a) = \begin{pmatrix} g(a) & 0 \\ 0 & g(\bar{a})^T \end{pmatrix} \text{ for } a \in \Gamma.$$

The list of matrices $(h(a); a \in \Gamma)$ can be computed in polynomial time and we have $h(\bar{a}) = \overline{h(a)}$. Now, for each regular language C_j , $1 \leq j \leq k$ we compute vectors $I_j, F_j \in \mathbb{B}^{2n}$ (corresponding to initial and final states) such that for all $w \in \Gamma^*$ and $1 \leq j \leq k$ we have the equivalence:

$$w \in C_j \Leftrightarrow I_j^T h(w) F_j = 1.$$

Having done these computations we make a non-deterministic guess $\rho(X) \in M_{2n}$ for each variable $X \in \Omega$. We verify $\rho(\bar{X}) = \overline{\rho(X)}$ for all $X \in \Omega$ and whenever there is a constraint of type $X \in C_j$ for some $1 \leq j \leq m$ (or $X \notin C_j$ for some $m < j \leq k$), then we verify $I_j^T \rho(X) F_j = 1$, if $1 \leq j \leq m$ (or $I_j^T \rho(X) F_j = 0$, if $m < j \leq k$).

This finishes the reduction of problem EFMI to EWC. \square

4 Problem Ewc is in Pspace

4.1 Road-Map

The proof of Theorem 8 is based on three transformation rules for equations with constraints. Each transformation preserves unsolvability; and it can be applied as long as the computation respects a given polynomial space bound. (The notion of *admissibility* given in Definition 31 formalizes the notion that the size of some object is bounded polynomially in the input size.)

No transformation rule introduces any new variable, but it may happen that the number of variables decreases. So, the global strategy is to apply the rules until all variables have been eliminated; the final step is then a direct evaluation of an equation without variables.

If the final output is *yes*, then the input equation is solvable, too. The main difficulty in the proof is the converse. We have to show that we can perform all these transformations within polynomial space such that for a solvable equation with constraints at least one computation path leads to the output *yes*.

In order to overcome this difficulty various notions and concepts are developed. We follow the approach of Plandowski [34, 35], but we have two sources for additional complications. We have to cope with the involution and we

have constraints. It is fairly standard to handle regular constraints. It may look rather technical if a reader sees it for the first time, but there is no surprise and the real additional difficulty is condensed in one subsection.

There are in fact three subsections 4.2, 4.6, and 4.9, where regular constraints play a crucial role. In 4.2 we show why an explicit specification of the constants is necessary. On an algebraic level we have to solve a membership problem in a submonoid of Boolean matrices. The submonoid is given by a list of matrices and we ask whether some other matrix A is a product of matrices from the list. Clearly, the answer may be *no*, but if we add A to the list, then it becomes trivially *yes*. In our language this means that it may happen that an equation with constraints becomes solvable by enlarging the alphabet of constants. This effect is not possible without constraints: If a word equation $L = R$ without constraints has a solution, then it has a solution over the alphabet of constants which appear in the string LR .

The presence of constraints makes it necessary to formalize the notion of *projection* in 4.6. A projection is a controlled way of introducing new constants such that unsolvable equations remain unsolvable. The use of new constants is inherent in Plandowski's method. If during the transformation the underlying word equations becomes too long, long subsequences of constants (factors) are coded as a single new letter. So, the alphabet of constants changes all the time: We remove constants in order to keep the alphabet size polynomially bounded, and introduce them in order to keep the length of the underlying word equation polynomially bounded. The technical preparation for this is done in Section 4.9. It introduces the notion of *free interval* and it is there where our presentation becomes more involved due to constraints.

Dealing with involutions is the main source for new difficulties. For example we cannot directly apply the usual method for bounding the exponent of periodicity. We need a new concept of p -stable normal form in Subsection 4.3. The result of this section is however as expected: If a w_0 represents a solution of minimal length, then the number of repetitions inside w_0 is bounded singly exponential in the size of the equation. Thus if $w_0 = uv^k w$, then in binary notation k uses polynomially many bits only.

This leads directly to Section 4.4. Word equations are not stored in plain form, but Plandowski's method uses data compression to keep them within polynomial size. More specifically, we allow regular expressions with exponents in binary notation.

The following three subsections explain the transformation rules in detail: In 4.5 we formalize the way to remove constants and 4.6 deals with the controlled way of introducing them. In 4.7 we formalize guessing a *partial solution*.

The transformation rules lead to the formal description of a *search graph*

in 4.8. The difficulty of proving Theorem 8 is reduced to showing that the search graph contains a path from a solvable input equation to some trivial equation. This part is very complex, but the basic ideas can be traced to [36] where Lempel-Ziv encodings of minimal solutions of a word equation are investigated. Key notions are *critical word* and ℓ -factorization. The technical part is developed in sections 4.10 to 4.15.

4.2 A PSPACE-Complete Subproblem

The following proposition states that two basic operations, which are used several times as subroutines, can be performed in PSPACE.

Proposition 13 *The following problems are PSPACE-complete with respect to the input size $n + \log |\Gamma|$.*

INPUT: A matrix $B \in \mathbb{B}^{n \times n}$ and a homomorphism $g : \Gamma^ \rightarrow \mathbb{B}^{n \times n}$ given as a list of matrices $(B_1, \dots, B_{|\Gamma|})$.*

QUESTION: Is there some $u \in \Sigma^$ such that $g(u) = B$?*

INPUT: A matrix $A \in M_{2n}$ and a morphism $h : \Gamma \rightarrow M_{2n}$ given as a list of matrices $(A_1, \dots, A_{|\Gamma|})$ with $\overline{A_{a_i}} = A_{\bar{a}_i}$ for all $a_i \in \Gamma$.

QUESTION: Is there some $w \in \Gamma^$ such that $h(w) = A$ and $w = \bar{w}$?*

Proof. The first problem is closely related to the intersection problem of regular languages and its PSPACE-hardness is again due to Kozen [22], see also [11, MS5]. The PSPACE-algorithm starts with the unit matrix. Then it guesses a word u letter by letter and, simultaneously, calculates $g(u)$: If we guess the letter a_i , then we move to the i -th matrix in the list $(B_1, \dots, B_{|\Gamma|})$ describing g , and we multiply B_i on the right to the current value held in the work space. We terminate if and only if $g(u) = B$.

The second problem can be solved since $w = \bar{w}$ implies $w = ub\bar{u}$ for some $u \in \Gamma^*$ and $b \in \Gamma \cup \{1\}$ with $b = \bar{b}$. Hence we can guess some B and b and we verify $A = Bh(b)\overline{B}$ and $b = \bar{b}$. Then using the first part, we check that $B = h(u)$ for some $u \in \Gamma^*$.

Since there is no reference which shows the PSPACE-hardness of the second problem, we sketch a reduction from the first to the second one: Consider a mapping $g : \Sigma \rightarrow \mathbb{B}^{n \times n}$ and $B \in \mathbb{B}^{n \times n}$, the pair (B, g) is an instance of the first problem. Let Γ be the disjoint union $\Sigma \cup \bar{\Sigma}$ and let $g(\bar{a}) = 1$, where $1 \in \mathbb{B}^{n \times n}$ is the identity matrix. In the notations of above we have $h(a) = \begin{pmatrix} g(a) & 0 \\ 0 & 1 \end{pmatrix}$ for $a \in \Gamma$. Let $A = \begin{pmatrix} B & 0 \\ 0 & B^T \end{pmatrix}$, then the pair (A, h) becomes an instance of the second problem. Clearly, if $g(u) = B$ for some

$u \in \Sigma^*$, then $h(u\bar{u}) = \begin{pmatrix} B & 0 \\ 0 & B^T \end{pmatrix}$. For the converse note that the matrices $h(a)$ and $h(\bar{b})$ commute for all $a, b \in \Sigma$. If there is some $w = \bar{w} \in \Gamma^*$ with $h(w) = \begin{pmatrix} B & 0 \\ 0 & B^T \end{pmatrix}$, then we can write $w = w_1\bar{w}_1$ and we may assume that $w_1 = u_1\bar{u}_2$ with $u_1, u_2 \in \Sigma^*$. It follows that $g(u_1u_2) = B$. \square

Assume that an equation $E = (\Gamma, h, \Omega, \rho; L = R)$ contains a variable X in the specification which does not occur in $LR\bar{L}\bar{R}$. In this case the equation might be unsolvable, simply because $\rho(X) \notin h(\Gamma^*)$. However, by Proposition 13 we can test this in PSPACE. Therefore, if X does not appear in $LR\bar{L}\bar{R}$ and $\rho(X) \in h(\Gamma^*)$, then we can remove X and \bar{X} from the specification. This yields the following remark.

Remark 14 *Henceforth, if $E = (\Gamma, h, \Omega, \rho; L = R)$ is an equation with constraints, then we assume that all variables occur somewhere in $LR\bar{L}\bar{R}$. As a consequence, we may assume $|\Omega| \leq 2|LR|$.*

4.3 The Exponent of Periodicity

A key step in proving Theorem 8 is to find an effective bound on the exponent of periodicity in a solution of minimal length. This idea is used in all known algorithms for solving word equations, c.f. [24, 34, 35]. It turns out that the well-known result on word equations [20] transfers to the situation here: The exponent of periodicity can be bounded by a singly exponential function. Let $w \in \Gamma^*$ be a word. The *exponent of periodicity* $\exp(w)$ is defined by

$$\exp(w) = \sup\{\alpha \in \mathbb{N} \mid \exists u, v, p \in \Gamma^*, p \neq 1 : w = up^\alpha v\}.$$

Proposition 15 *Let $E = (\Gamma, h, \Omega, \rho; L = R)$ be a solvable equation with constraints. Then there is a solution $\sigma : \Omega \rightarrow \Gamma^*$ such that $\exp(\sigma(L)) \in 2^{\mathcal{O}(d+n \log n)}$.*

The proof of Proposition 15 is independent of the rest of the paper. Therefore we postpone it to the appendix, Section 6.

4.4 Exponential Expressions

In order to keep the computation in polynomial space Plandowski's method uses *exponential expressions*. We give inductive definitions for an exponential expression, its evaluation, and its size.

Definition 16 • Every word $w \in \Gamma^*$ is an exponential expression. The evaluation $\text{eval}(w)$ is equal to w , its size $\|w\|$ is equal to the length $|w|$.

- Let e, e' be exponential expressions. Then ee' is an exponential expression. Its evaluation is the concatenation $\text{eval}(ee') = \text{eval}(e)\text{eval}(e')$, its size is $\|ee'\| = \|e\| + \|e'\|$.
- Let e be an exponential expression and $k \in \mathbb{N}$. Then $(e)^k$ is an exponential expression. Its evaluation is $\text{eval}((e)^k) = (\text{eval}(e))^k$, its size is $\|(e)^k\| = \|e\| + \max\{1, \lceil \log_2(k) \rceil\}$.

According to a standard notion in combinatorics on words a *factor* of a word w is a word u such that $w = w_1uw_2$. A factor u is called *proper* if $1 \neq u \neq w$.

Lemma 17 Let $u \in \Gamma^*$ be a factor of a word $w \in \Gamma^*$. Assume that w can be represented by some exponential expression of size p . Then we can find an exponential expression of size at most p^2 that represents u .

Proof. The proof is an easy argument by structural induction. □

Lemma 17 will be applied to exponential expressions where the size $\|e\|$ is bounded by some value which is polynomial in the input size of the equation E_0 . Since the size of the exponential expressions for factors can be the square of the original polynomial, we can apply this subroutine in nested way a constant number of times, only. In our application the nested depth does not go beyond two.

The next lemma is straightforward since we allow a polynomial space bound without any time restriction. The proof of Lemma 18 is omitted.

Lemma 18 The following two problems can be solved in PSPACE.

INPUT: Exponential expressions e and e' .

QUESTION: Do we have $\text{eval}(e) = \text{eval}(e')$?

INPUT: A mapping $h : \Gamma \rightarrow M_{2n}$ and an exponential expression e .

OUTPUT: The matrix $h(\text{eval}(e)) \in M_{2n}$.

Remark 19 The computation above can actually be performed in polynomial time, but this is not evident for the first question, see [32] for details.

Henceforth, we allow that the part $L = R$ of an equation with constraints may be given by a pair of exponential expressions (e_L, e_R) with $\text{eval}(e_L) = L$ and $\text{eval}(e_R) = R$.

Definition 20 Let $E = (\Gamma, h, \Omega, \rho; e_L = e_R)$ and $E' = (\Gamma, h, \Omega, \rho; e'_L = e'_R)$ be equations with constraints. We write $E \equiv E'$, if $\text{eval}(e_L) = \text{eval}(e'_L)$ and $\text{eval}(e_R) = \text{eval}(e'_R)$ as strings in $(\Gamma \cup \Omega)^*$.

The meaning of $E \equiv E'$ is that E and E' represent exactly the same equation if they were written out explicitly. By Lemma 18 we can decide $E \equiv E'$ in polynomial space; moreover, Remark 19 says that this decision is actually possible in polynomial time.

4.5 Base Changes

In this subsection we describe the first transformation rule. Let $h : \Gamma^* \rightarrow M_{2n}$ be a morphism. Let $(\Gamma', \bar{\cdot})$ be another alphabet with involution and let $\beta : \Gamma' \rightarrow \Gamma^*$ be some mapping such that $\beta(\bar{a}) = \overline{\beta(a)}$ for all $a \in \Gamma'$. We define $h' : \Gamma' \rightarrow M_{2n}$ by $h' = h\beta$. We extend β to a morphism $\beta : (\Gamma' \cup \Omega)^* \rightarrow (\Gamma \cup \Omega)^*$ by leaving the variables invariant and we call the morphism β a *base change*. Let β be a base change and $E' = (\Gamma', h\beta, \Omega, \rho; L' = R')$ be an equation with constraints. The equation $\beta_*(E')$ is defined by

$$\beta_*(E') = (\Gamma, h, \Omega, \rho; \beta(L') = \beta(R')).$$

Lemma 21 Let E' be an equation with constraints and $\beta : \Gamma' \rightarrow \Gamma^*$ be a base change. If σ' is a solution of E' , then $\sigma = \beta\sigma'$ is a solution of $\beta_*(E')$.

Proof. Clearly $\sigma(\bar{X}) = \overline{\sigma(X)}$ and $h\sigma(X) = h\beta\sigma'(X) = h'\sigma'(X) = \rho(X)$ for all $X \in \Omega$. Next by definition $\sigma(a) = a$ for $a \in \Gamma$ and $\beta(X) = X$ for $X \in \Omega$. Hence $\sigma\beta(a) = \beta\sigma'(a)$ for $a \in \Gamma'$ and therefore $\sigma\beta = \beta\sigma' : (\Gamma' \cup \Omega)^* \rightarrow \Gamma^*$. This means $\sigma\beta(L) = \beta\sigma'(L) = \beta\sigma'(R) = \sigma\beta(R)$ since $\sigma'(L) = \sigma'(R)$. \square

Lemma 21 leads to the first rule.

Rule 1 If we have $E \equiv \beta_*(E')$ and we are looking for a solution of E , then it is enough to find a solution for E' . Hence, during a non-deterministic search we may replace E by E' .

For readability of the following examples all constraints are defined by membership in a regular language rather than by a mapping ρ . We also strengthen constraints in the examples (thereby having fewer solutions) in order to avoid lengthy regular expressions.

Example 22 Let $\Gamma = \{a, b, c, \bar{a}, \bar{b}, \bar{c}\}$. Consider the following equation E :

$$X\bar{X} = Y\bar{b}\bar{c}\bar{b}\bar{a}\bar{b}\bar{c}\bar{b}Y\bar{Z}abc\bar{b}\bar{Y}$$

with constraints $X \in \Gamma^{300}\Gamma^*$ and $Z \in \bar{b}\bar{c}\bar{b}\bar{a}\Gamma^*$. Define $\Gamma' = \{a, b, \bar{a}, \bar{b}\}$ and a base change $\beta : \Gamma' \rightarrow \Gamma^*$ by $\beta(a) = abcb$ and $\beta(b) = bcb$. Then the equation E is of the form $\beta_*(E')$ where E' is given by

$$X\bar{X} = Y\bar{a}\bar{b}Y Z a\bar{Y}.$$

We may strengthen the constraint to $X \in \Gamma'^{100}\Gamma'^*$ and $Z \in \bar{a}\Gamma'^*$. According to Rule 1 it is enough to solve E' . The effect of the base change β is that both the equation E' and the alphabet of constants are smaller. (The letter c is not used anymore.) Note also that the length restriction on X switched from $|X| \geq 300$ to $|X| \geq 100$. However, base changes may have a prize: It might be that $E = \beta_*(E')$ has a solution, whereas E' is unsolvable. As we will see later, in our example the guess has been correct in the sense that E' has a solution.

4.6 Projections

Let $(\Gamma, \bar{\quad})$ and $(\Gamma', \bar{\quad})$ be alphabets with involution such that $\Gamma \subseteq \Gamma'$. A *projection* is a morphism $\pi : \Gamma'^* \rightarrow \Gamma^*$ such that both, $\pi(a) = a$ for $a \in \Gamma$ and $\pi(\bar{a}) = \overline{\pi(a)}$ for all $a \in \Gamma'$.

Let E be an equation with constraints $E = (\Gamma, h, \Omega, \rho; L = R)$. Then we define an equation with constraints $\pi^*(E)$ by

$$\pi^*(E) = (\Gamma', h\pi, \Omega, \rho; L = R).$$

The equation $\pi^*(E)$ uses a larger alphabet of constants than E does, but the word equation $L = R$ is exactly the same. Therefore $\pi^*(E)$ uses constants which do not appear in $L = R$. These constants may help to find (short) solutions which satisfy regular constraints. Note that every projection $\pi : \Gamma'^* \rightarrow \Gamma^*$ defines a base change π such that $\pi_*\pi^*(E) = E$. Let $E' = \pi^*(E)$. By Rule 1 we may replace $\pi_*(E')$ by $\pi^*(E)$. We formulate this special case a second rule.

Rule 2 *Let π be a projection. If we are looking for a solution of E , then it is enough to find a solution for $\pi^*(E)$. Hence, during a non-deterministic search we may replace E by $\pi^*(E)$.*

Remark 23 *The reason to introduce Rule 2 will become clear only later. In Section 4.8 we define the formal notion of a search graph. We restrict the use of Rule 1 to so-called admissible base changes (c.f. Definition 31), whereas there is no such restriction for the projection π when we apply Rule 2.*

Lemma 24 *Let $E = (\Gamma, h, \Omega, \rho; L = R)$ and $E' = (\Gamma', h', \Omega, \rho; L = R)$ be equations with constraints. Then the following two statements hold.*

- i) There is a projection $\pi : \Gamma'^* \rightarrow \Gamma^*$ such that $\pi^*(E) = E'$, if and only if both, $h'(\Gamma') \subseteq h(\Gamma^*)$ and for all $a \in \Gamma'$ with $a = \bar{a}$ there is some $w \in \Gamma^*$ with $w = \bar{w}$ such that $h'(a) = h(w)$.*
- ii) Let $\pi^*(E) = E'$ for some projection π and let $\sigma' : \Omega \rightarrow \Gamma'^*$ be a solution of E' . Then there is a solution σ for E such that $|\sigma(L)| \leq 2|M_{2n}||\sigma'(L)|$.*

Proof. i) Clearly, the only-if condition is satisfied by the definition of a projection since then $h' = h\pi$. For the converse, assume that $h'(\Gamma') \subseteq h(\Gamma^*)$ and that $a = \bar{a}$ implies $h'(a) \in h(\{w \in \Gamma^* \mid w = \bar{w}\})$. Then for each $a \in \Gamma' \setminus \Gamma$ we can choose a word $w_a \in \Gamma^*$ such that $h'(a) = h(w_a)$. We can make the choice such that $w_{\bar{a}} = \overline{w_a}$ for all $a \in \Gamma' \setminus \Gamma$. If $a \neq \bar{a}$, then we can find w_a such that $|w_a| < |M_{2n}|$, since we can take the shortest word $w_a \in \Gamma^*$ such that $h(w_a) = h'(a) \in M_{2n}$. For $a = \bar{a}$ we know that there is some word $w_a \in \Gamma^*$ with $h'(a) = h(w_a)$ and $w_a = \overline{w_a}$. Hence we can write $w_a = vb\bar{v}$ with $b \in \Gamma \cup \{1\}$ and $b = \bar{b}$. For $b \neq 1$ we can demand $|w_a| \leq 2|M_{2n}| - 1$. For $b = 1$ we can demand $|w_a| \leq 2|M_{2n}| - 2$. Thus, we find a projection $\pi : \Gamma'^* \rightarrow \Gamma^*$ such that $\pi^*(E) = E'$ and moreover, $|\pi(a)| < 2|M_{2n}|$ for all $a \in \Gamma'$.

ii) According to the proof of i) we may assume that $\pi : \Gamma'^* \rightarrow \Gamma^*$ satisfies $|\pi(a)| < 2|M_{2n}|$ for all $a \in \Gamma'$. Since π defines a base change with $\pi_*(E') = E$, we know by Lemma 21 that $\sigma = \pi\sigma'$ is a solution of E . Clearly, $|\sigma(L)| = |\pi\sigma'(L)| \leq 2|M_{2n}||\sigma'(L)|$. \square

Example 25 *Let us continue with the equation which has been obtained by the transformation in Example 22. In order to simplify notations, we let E be the equation $X\bar{X} = Y\bar{a}\bar{b}Y Z\bar{a}\bar{Y}$, and $\Gamma = \{a, b, \bar{a}, \bar{b}\}$.*

Remember that the constraint on X has changed to $|X| \geq 100$. Let us reintroduce a letter c and put $\Gamma' = \{a, b, c, \bar{a}, \bar{b}, \bar{c}\}$. We may define a projection $\pi : \Gamma' \rightarrow \Gamma^$ by $\pi(c) = b^{100}$. The equation $E' = \pi^*(E)$ looks as above, but in E' the constraint for X has changed. The new constraint is $|X| \geq 100$ or $X \in \Gamma^*c\Gamma^*$. Thus, a solution for X might be very short now.*

During the procedure we occasionally have to decide whether there is a projection $\pi : \Gamma'^* \rightarrow \Gamma^*$ such that $\pi^*(E) = E'$. This decision is possible according to the next proposition.

Proposition 26 *The following problem PSPACE-complete.*

INPUT: Alphabets $(\Gamma, \bar{\cdot}) \subseteq (\Gamma', \bar{\cdot})$ and mappings h, h' , where h is the restriction of h' .

QUESTION: Is there a projection $\pi : \Gamma'^* \rightarrow \Gamma^*$ such that $h' = h\pi$?

Proof. This follows from Lemma 24 i) and Proposition 13. \square

4.7 Partial Solutions

Let $\Omega' \subseteq \Omega$ be a subset of the variables which is closed under involution. We assume that there is a mapping $\rho' : \Omega' \rightarrow M_{2n}$ with $\rho'(\overline{X}) = \overline{\rho'(X)}$, but we do not require that ρ' is the restriction of $\rho : \Omega \rightarrow M_{2n}$. Consider an equation with constraints $E = (\Gamma, h, \Omega, \rho; e_L = e_R)$. A *partial solution* is a mapping $\delta : \Omega \rightarrow \Gamma^* \Omega' \Gamma^* \cup \Gamma^*$ such that the following conditions are satisfied:

- i) $\delta(X) \in \Gamma^* X \Gamma^*$ for all $X \in \Omega'$,
- ii) $\delta(X) \in \Gamma^*$ for all $X \in \Omega \setminus \Omega'$,
- iii) $\delta(\overline{X}) = \overline{\delta(X)}$ for all $X \in \Omega$.

The mapping δ is extended to a morphism $\delta : (\Gamma \cup \Omega)^* \rightarrow (\Gamma \cup \Omega')^*$ by leaving the elements of Γ invariant. Let $E' = (\Gamma, h, \Omega', \rho'; e_{L'} = e_{R'})$ be another equation with constraints (using the same Γ and h). By abuse of language, we write $E' \equiv \delta_*(E)$, if there exists some partial solution $\delta : \Omega \rightarrow \Gamma^* \Omega' \Gamma^* \cup \Gamma^*$ such that the following conditions hold: $L' = \delta(L)$, $R' = \delta(R)$, $\rho(X) = h(u)\rho'(X)h(v)$ for $\delta(X) = uXv$, and $\rho(X) = h(w)$ for $\delta(X) = w \in \Gamma^*$.

Lemma 27 *In the notation of above, let $E' \equiv \delta_*(E)$ for some partial solution $\delta : \Omega \rightarrow \Gamma^* \Omega' \Gamma^* \cup \Gamma^*$. If σ' is a solution of E' , then $\sigma = \sigma'\delta$ is a solution of E . Moreover, we have $\sigma(L) = \sigma'(L')$ and $\sigma(R) = \sigma'(R')$.*

Proof. By definition, δ and σ' are extended to morphisms $\delta : (\Gamma \cup \Omega)^* \rightarrow (\Gamma \cup \Omega')^*$ and $\sigma' : (\Gamma \cup \Omega')^* \rightarrow \Gamma^*$ leaving the letters of Γ invariant. Since $E' = \delta_*(E)$ we have $\delta(L) = L'$ and $\delta(R) = R'$. Since σ' is a solution, we have $\sigma(L) = \sigma'\delta(L) = \sigma'(L') = \sigma'(R') = \sigma'\delta(R) = \sigma(R)$ and σ leaves the letters of Γ invariant. The solution σ' satisfies $h\sigma'(X) = \rho'(X)$ for all $X \in \Omega'$. Hence, if $\delta(X) = uXv$, then $\rho(X) = h(u)\rho'(X)h(v) = h(u\sigma'(X)v) = h\sigma'(uXv) = h\sigma'\delta(X) = h\sigma(X)$. If $\delta(X) = w \in \Gamma^*$, then $\sigma(X) = \sigma'\delta(X) = w$ and $\rho(X) = h(w)$, again by the definition of a partial solution. \square

Lemma 28 *The following problem can be solved in PSPACE.*

INPUT: Two equations with constraints $E = (\Gamma, h, \Omega, \rho; e_L = e_R)$ and $E' = (\Gamma, h, \Omega', \rho'; e_{L'} = e_{R'})$.

QUESTION: Is there some partial solution δ such that $\delta_*(E) \equiv E'$?

If $\delta_*(E) \equiv E'$ is true, then there are exponential expressions of polynomial size e_u, e_v for each $X \in \Omega'$ and e_w for each $X \in \Omega \setminus \Omega'$ such that

$$\begin{aligned} \delta(X) &= \text{eval}(e_u)X\text{eval}(e_v) && \text{for } X \in \Omega', \\ \delta(X) &= \text{eval}(e_w) && \text{for } X \in \Omega \setminus \Omega'. \end{aligned}$$

Proof. Let $L = \text{eval}(e_L)$, $R = \text{eval}(e_R)$, $L' = \text{eval}(e_{L'})$, and $R' = \text{eval}(e_{R'})$. The non-deterministic PSPACE algorithm works as follows:

For each variable $X \in \Omega'$ we guess exponential expressions e_u and e_v with $\text{eval}(e_u), \text{eval}(e_v) \in \Gamma^*$. We define exponential expressions $e_X = e_u X e_v$ and we define $\delta(X) = \text{eval}(e_X)$. For each $X \in \Omega \setminus \Omega'$ we guess an exponential expression e_X with $\text{eval}(e_X) \in \Gamma^*$ and we define $\delta(X) = \text{eval}(e_X)$.

Next we verify whether or not $\delta_*(E) \equiv E'$. During this test we have to create an exponential expression f_L (and f_R , resp.) by replacing X in e_L (and e_R , resp.) with the expression e_X . This increases the size in the worst case by a factor of $\max\{||e_X|| \mid X \in \Omega\}$. The other tests whether $\rho(X) = h(u)\rho'(X)h(v)$ for $\delta(X) = uXv$ and $\rho(X) = h(w)$ for $\delta(X) = w \in \Gamma^*$ involve exponential expressions over Boolean matrices and can be done in polynomial time.

The correctness of the algorithm follows from our general assumption (Remark 14) that all $X \in \Omega$ appear in $LR\overline{LR}$. Therefore, if we have $\delta_*(E) \equiv E'$, then every factor of $\delta(X)$ (or of $\delta(\overline{X})$) appears necessarily as a factor in $L'R' = \delta(LR)$. Hence every factor of $\delta(X)$ has an exponential expression of polynomial size by Lemma 17. \square

Remark 29 *Actually, the test for $\delta_*(E) \equiv E'$ can be performed in non-deterministic polynomial time by Remark 19.*

Lemma 27 leads to the third and last rule.

Rule 3 *If δ is a partial solution and if we are looking for a solution of E , then it is enough to find a solution for $\delta_*(E)$. Hence, during a non-deterministic search we may replace E by $\delta_*(E)$.*

Example 30 *We continue with our running example. After renaming, the equation E is given by*

$$X\overline{X} = Y\overline{a}bY Z a\overline{Y},$$

and the alphabet of constant is given by $\Gamma = \{a, b, c, \bar{a}, \bar{b}, \bar{c}\}$. We strengthen constraints such that $X \in \Gamma^*c\Gamma^*$ and $Z \in \bar{a}\{a, b, \bar{a}, \bar{b}\}^*$.

We may guess the partial solution as follows: $\delta(X) = aX$, $\delta(Y) = Y$, and $\delta(Z) = \bar{a}b$. The new equation $\delta_*(E)$ is

$$aX\bar{X}\bar{a} = Y\bar{a}bY\bar{a}b\bar{Y}.$$

The remaining constraint is that the solution for X has to use the letter c . The process can continue, for example, we can apply Rule 1 again by defining another base change $\beta(b) = ba$ to get the equation

$$aX\bar{X}\bar{a} = Y\bar{b}Y\bar{a}b\bar{Y}$$

over $\Gamma = \{a, b, c, \bar{a}, \bar{b}, \bar{c}\}$. Since the last equation has a solution (e.g., given by $\sigma(X) = bc\bar{c}bbabc$ and $\sigma(Y) = abc\bar{c}b$), the first equation with constraints in Example 22 has a solution too.

4.8 The Search Graph and Plandowski's Algorithm

The input of Problem EWC is an equation with constraints. In order to fix notations we call it $E_0 = (\Gamma_0, h_0, \Omega_0, \rho_0; L_0 = R_0)$ and we let $d = |L_0R_0|$. According to Remark 14 we assume $|\Omega_0| \leq 2d$.

Definition 31 Let p_0 be a polynomial. The notion of admissibility is defined with respect to $p_0(\|E_0\|)$.

- An exponential expression e is admissible, if $\|e\| \leq p_0(\|E_0\|)$.
- A base change $\beta : \Gamma' \rightarrow \Gamma^*$ is admissible, if $|\Gamma'| \leq p_0(\|E_0\|)$ and for all $a \in \Gamma'$ there is an admissible exponential expression for $\beta(a)$.
- An equation with constraints $E = (\Gamma, h, \Omega, \rho; e_L = e_R)$ is admissible, if $|\Gamma \setminus \Gamma_0| \leq p_0(\|E_0\|)$, $h(a) = h_0(a)$ for $a \in \Gamma_0$, and $e_L e_R$ is admissible.

In the following we assume that a polynomial p_0 (of large enough degree) has been fixed whenever we speak about admissibility. We do not calculate p_0 explicitly, but it will become clear from the context what *large enough* actually means.

Definition 32 The search graph of E_0 is a directed graph where nodes are admissible equations with constraints. For two nodes E, E' there is an arc $E \rightarrow E'$, if there are an admissible base change β , a projection π , and a partial solution δ such that $\delta_*(\pi^*(E)) \equiv \beta_*(E')$.

Lemma 33 *Let p_0 be a polynomial of degree at least 1. The following problem is PSPACE-complete.*

INPUT: Equations with constraints E_0 , E , and E' such that E and E' are admissible with respect to $p_0(\|E_0\|)$.

QUESTION: Is there an arc $E \rightarrow E'$ in the search graph of E_0 ?

Proof. The arc from E to E' is established by applying Rules 2, 1, and 3 (in this order) to E . More precisely, we let $E_0 = (\Gamma_0, h_0, \Omega_0, \rho_0; L_0 = R_0)$, $E = (\Gamma, \Omega, \rho; e_L = e_R)$, and $E' = (\Gamma', h', \Omega', \rho'; e_{L'} = e_{R'})$. We first guess some alphabet $(\Gamma'', \bar{\cdot})$ of polynomial size together with $h'' : \Gamma'' \rightarrow M_{2n}$ and we guess some admissible base change $\beta : \Gamma' \rightarrow \Gamma''^*$ such that $h' = h''\beta$. We compute $\beta_*(E')$. Next, we guess some equation with constraints E'' which uses Γ'' and Ω . We check using Lemma 28 that there is some partial solution $\delta : \Omega \rightarrow \Gamma''^* \Omega' \Gamma''^* \cup \Gamma''^*$ such that $\delta_*(E'') \equiv \beta_*(E')$. (Note that every equation with constraints E'' satisfying $\delta_*(E'') \equiv \beta_*(E')$ for some δ can be represented in polynomial space by Lemma 17.) Finally we check using Proposition 26 that there is some projection $\pi : \Gamma'' \rightarrow \Gamma$ such that $\pi^*(E) \equiv E''$. We obtain $\delta_*(\pi^*(E)) \equiv \beta_*(E')$.

The PSPACE-hardness follows by Proposition 13 which shows that the problem is PSPACE-hard on instances of the following type: The equation for E and E' is $X = X$, we have $\rho(X) = \rho'(X) = A \in M_{2n}$, and $\Gamma' \setminus \Gamma = \{a, \bar{a}\}$ with $h'(a) = A$. \square

Remark 34 *Following Remarks 19 and 29 the problem in Lemma 33 can be decided in non-deterministic polynomial time, if the monoid M_{2n} is not part of the input and the parameter n is viewed as a constant.*

On a high-level description, Plandowski's algorithm applies Rules 1 to 3 in a non-deterministic way until a trivial equation is found. An actual implementation of the algorithm depends on the chosen polynomial p_0 and it has the following structure.

begin

$E := E_0$

while $\Omega \neq \emptyset$ **do**

Guess an equation with constraints E' ,

which is admissible with respect to $p_0(\|E_0\|)$

Verify that $E \rightarrow E'$ is an arc in the search graph of E_0

$E := E'$

endwhile

return “eval(e_L) = eval(e_R)”

end

Lemmata 21, 24 ii), and 27 say that the algorithm returns *true* only if E_0 is solvable. The proof of Theorem 8 is therefore reduced to the statement that there is a polynomial p_0 such that for all E_0 we have, if E_0 is solvable, then the search graph contains a path to some solvable equation without variables.

Remark 35 *If the arc $E \rightarrow E'$ is due to some $\pi : \Gamma''^* \rightarrow \Gamma^*$, $\delta : \Omega \rightarrow \Gamma''^* \Omega' \Gamma''^* \cup \Gamma''^*$, and $\beta : \Gamma'^* \rightarrow \Gamma''^*$, then a solution $\sigma' : \Omega' \rightarrow \Gamma'^*$ of E' yields the solution $\sigma = \pi(\beta\sigma')\delta$. Hence we may assume that the length of a solution has increased by at most an exponential factor by Lemma 24 ii). Since we are going to perform the search in a graph of at most exponential size, we automatically get a doubly exponential upper bound for the length of a minimal solution by backwards computation on such a path. This is still the best known upper bound (although a singly exponential bound is conjectured), see [33].*

4.9 Free Intervals

For a word $w \in \Gamma^*$ we let $\{0, \dots, |w|\}$ be the set of its *positions*. The idea is that factors of w are between positions. To be more specific, let $w = a_1 \cdots a_m$ be a word with $a_i \in \Gamma$. Then $[\alpha, \beta]$ with $0 \leq \alpha < \beta \leq m$ is called a *positive interval* and the word $w[\alpha, \beta]$ is defined as the factor $a_{\alpha+1} \cdots a_\beta$.

It is convenient to have an involution on the set of intervals. If $[\alpha, \beta]$ is a positive interval, then $[\beta, \alpha]$ is also called a (non-positive) interval, and we define $w[\beta, \alpha] = \overline{w[\alpha, \beta]}$. Moreover, we define $w[\alpha, \alpha]$ to be the empty word. For all $0 \leq \alpha, \beta \leq m$ we let $\overline{[\alpha, \beta]} = [\beta, \alpha]$; therefore, $w[\alpha, \beta] = \overline{w[\beta, \alpha]}$.

In the following we assume that the input equation E_0 has a solution σ with $w_0 = \sigma(L_0) = \sigma(R_0)$ and $m_0 = |w_0|$. We have $w_0 \in \Gamma_0^*$, but in this section the alphabet Γ_0 is replaced by some other alphabet Γ , which turns out to be a set of non-empty words over Γ_0 . We let d be the denotational length of the equation and $L_0 = x_1 \cdots x_g$ and $R_0 = x_{g+1} \cdots x_d$, $x_i \in (\Gamma_0 \cup \Omega_0)$ for $1 \leq i \leq d$. We assume $2 \leq g < d < m_0$ whenever necessary. We also make the assumption that $\sigma(x_i) \neq 1$ for all $1 \leq i \leq d$. This assumption can be realized e.g. in the preprocessing.

We are going to define an equivalence relation \approx on the set of intervals of w_0 . For this we need some preparation. For $i \in \{1, \dots, d\}$ we define positions $l(i)$ and $r(i)$ such that $\sigma(x_i)$ starts in w_0 at the left position $l(i)$ and it ends at the right position $r(i)$. Formally, we define $l(i) \in \{0, \dots, m_0 - 1\}$ and $r(i) \in \{1, \dots, m_0\}$ by the congruences:

$$\begin{aligned} l(i) &\equiv |\sigma(x_1 \cdots x_{i-1})| \bmod m_0 \\ r(i) &\equiv |\sigma(x_1 \cdots x_i)| \bmod m_0 \end{aligned}$$

The next lemma says that subintervals of free intervals are free again.

Lemma 38 *Let $[\alpha, \beta]$ be a free interval and μ, ν such that $\min\{\alpha, \beta\} \leq \mu, \nu \leq \max\{\alpha, \beta\}$. Then the interval $[\mu, \nu]$ is also free.*

Proof. We may assume that $\alpha \leq \mu < \nu \leq \beta$. By contradiction assume that $[\mu, \nu]$ is not free. Then there is some $k \geq 0$ and some cut γ' such that

$$[\mu, \nu] = [\mu_0, \nu_0] \sim [\mu_1, \nu_1] \sim \cdots \sim [\mu_k, \nu_k]$$

with $\min\{\mu_k, \nu_k\} < \gamma' < \max\{\mu_k, \nu_k\}$. If $k = 0$, then we have an immediate contradiction. For $k \geq 1$ the relation $[\mu, \nu] \sim [\mu_1, \nu_1]$ is due to some pair x_i, x_j with $x_i = x_j$ or $x_i = \bar{x}_j$. Since $[\alpha, \beta]$ contains no cut, the same pair x_i, x_j defines an interval $[\alpha_1, \beta_1]$ such that $[\alpha, \beta] \sim [\alpha_1, \beta_1]$ and $\min\{\alpha_1, \beta_1\} \leq \mu_1, \nu_1 \leq \max\{\alpha_1, \beta_1\}$. Using induction on k we see that $[\alpha_1, \beta_1]$ is not free. But then $[\alpha, \beta]$ is not free, and this is a contradiction. \square

Next we introduce the notion of *implicit cut* for non-free intervals. For our purpose it is enough to define it for positive intervals. So, let $0 \leq \alpha < \beta \leq m_0$ such that $[\alpha, \beta]$ is not free. A position γ with $\alpha < \gamma < \beta$ is called an *implicit cut* of $[\alpha, \beta]$, if there is a cut γ' and an interval $[\alpha', \beta']$ such that

$$\begin{aligned} \min\{\alpha', \beta'\} &< \gamma' < \max\{\alpha', \beta'\}, \\ [\alpha, \beta] &\approx [\alpha', \beta'], \\ \gamma - \alpha &= |\gamma' - \alpha'|. \end{aligned}$$

The following observation will be used throughout. If we have $\alpha \leq \mu < \gamma < \nu \leq \beta$ and γ is an implicit cut of $[\alpha, \beta]$, then γ is also an implicit cut of $[\mu, \nu]$. In particular, neither $[\mu, \nu]$ nor $[\nu, \mu]$ is a free interval.

Definition 39 *A free interval $[\alpha, \beta]$ is called maximal free, if there is no free interval $[\alpha', \beta']$ such that both, $\alpha' \leq \min\{\alpha, \beta\} \leq \max\{\alpha', \beta'\} \leq \beta'$ and $|\beta - \alpha| < \beta' - \alpha'$.*

Lemma 40 states that maximal free intervals do not overlap.

Lemma 40 *Let $0 \leq \alpha \leq \alpha' < \beta \leq \beta' \leq m_0$ such that $[\alpha, \beta]$ and $[\alpha', \beta']$ are free intervals. Then the interval $[\alpha, \beta']$ is free, too.*

Proof. Assume by contradiction that $[\alpha, \beta']$ is not free. Then it contains an implicit cut γ with $\alpha < \gamma < \beta'$. By the observation above: If $\gamma < \beta$, then γ is an implicit cut of $[\alpha, \beta]$ and $[\alpha, \beta]$ is not free. Otherwise, $\alpha' < \gamma$ and $[\alpha', \beta']$ is not free. \square

Lemma 41 states the main observation of this section.

Lemma 41 *Let $[\alpha, \beta]$ be a maximal free interval. Then there are intervals $[\gamma, \delta]$ and $[\gamma', \delta']$ such that $[\alpha, \beta] \approx [\gamma, \delta] \approx [\gamma', \delta']$ and γ and δ' are cuts.*

Proof. We may assume that $\alpha < \beta$. We show the existence of $[\gamma, \delta]$ where $[\alpha, \beta] \approx [\gamma, \delta]$ and γ is a cut. (The existence of $[\gamma', \delta']$ where $[\alpha, \beta] \approx [\gamma', \delta']$ and δ' is a cut follows by a symmetric argument.)

If $\alpha = 0$, then α is a cut and we can choose $\delta = \beta$. Hence let $1 \leq \alpha$ and consider the positive interval $[\alpha - 1, \beta]$. This interval is not free and the only possible position for an implicit cut is α . Thus, for some cut γ we have $[\alpha - 1, \beta] \approx [\alpha', \beta']$ with $\min\{\alpha', \beta'\} < \gamma < \max\{\alpha', \beta'\}$ and $|\gamma - \alpha'| = 1$. A simple reflection shows that we have $[\alpha - 1, \alpha] \approx [\alpha', \gamma]$ and $[\alpha, \beta] \approx [\gamma, \beta']$. Hence we can choose $\delta = \beta'$. \square

In the following proposition the symbol Γ refers to some set of factors of the word w_0 . (Recall that $w_0 = \sigma(L_0) = \sigma(R_0)$ and σ is a solution of the input equation E_0 .) The set Γ becomes the basic alphabet later.

Proposition 42 *Let Γ be the set of words $w \in \Gamma_0^*$ such that there is a maximal free interval $[\alpha, \beta]$ with $w = w_0[\alpha, \beta]$. Then Γ is a subset of Γ_0^+ of size at most $2d - 2$. The set Γ is closed under involution.*

Proof. Let $[\alpha, \beta]$ be maximal free. Then $|\beta - \alpha| \geq 1$ and $[\beta, \alpha]$ is also maximal free by definition. Hence $\Gamma \subseteq \Gamma_0^+$ and Γ is closed under involution. By Lemma 41 we may assume that α is a cut. Say $\alpha < \beta$. Then $\alpha \neq m_0$ and there is no other maximal free interval $[\alpha, \beta']$ with $\alpha < \beta'$ because of Lemma 40. Hence there are at most $d - 1$ such intervals $[\alpha, \beta]$. Symmetrically, there are at most $d - 1$ maximal free intervals $[\alpha, \beta]$ where $\beta < \alpha$ and α is a cut. \square

For the moment let $\Gamma'_0 = \Gamma_0 \cup \Gamma$ where $\Gamma \subseteq \Gamma_0^+$ is the set defined in Proposition 42. The inclusion $\Gamma'_0 \subseteq \Gamma_0^+$ defines a natural projection $\pi : \Gamma'_0 \rightarrow \Gamma_0^*$ and a mapping $h'_0 : \Gamma'_0 \rightarrow M_{2n}$ by $h'_0 = h_0\pi$. Consider the equation with constraints $\pi^*(E_0)$, this is a node in the search graph, because the size of Γ is linear in d .

The reason to switch from Γ_0 to Γ'_0 is that, due to the constraints, the word w_0 may have long free intervals, even in a minimal solution. Over Γ'_0 long free intervals can be avoided. Formally, we replace w_0 by a solution w'_0 where $w'_0 \in \Gamma^*$. The definition of w'_0 is based on a factorization of w_0 into maximal free intervals. There is a unique sequence $0 = \alpha_0 < \alpha_1 < \dots < \alpha_k = m_0$ such that $[\alpha_{i-1}, \alpha_i]$ is a maximal free interval for each $1 \leq i \leq k$ and

$$w_0 = w_0[\alpha_0, \alpha_1] \cdots w_0[\alpha_{k-1}, \alpha_k].$$

Note that all cuts occur as some α_p , therefore we can think of the factors $w_0[\alpha_{i-1}, \alpha_i]$ as letters in Γ for $1 \leq i \leq k$. Moreover, all constants which appear in $L_0 R_0$ are elements of Γ , too. We replace w_0 by the word $w'_0 \in \Gamma^*$. Then we can define $\sigma' : \Omega_0 \rightarrow \Gamma^*$ such that both, $\sigma'(L_0) = \sigma'(R_0) = w'_0$ and $\rho_0 = h'_0 \sigma'$. In other words, σ' is a solution of $\pi^*(E_0)$. We have $w_0 = \pi(w'_0)$ and $\exp(w'_0) \leq \exp(w_0)$. The crucial point is that w'_0 has no long free intervals anymore. With respect to w'_0 and Γ'_0 all maximal free intervals have length exactly one.

Example 43 *Following Example 37, we use the same equation $aX\bar{X}\bar{a} = Y\bar{b}Y\bar{a}b\bar{Y}$ and we consider the solution w_0 .*

The new solution is defined by replacing in w_0 each factor bc by a new letter d which represents a maximal free interval. The new w_0 has the form

$$w_0 = \overset{0}{|} \overset{1}{a} \overset{\bar{3}}{|} \overset{\bar{4}}{d\bar{d}} \overset{\bar{6}}{|} \overset{\bar{7}}{b} \overset{\bar{8}}{|} \overset{\bar{9}}{ad} \overset{\bar{11}}{|} \overset{\bar{12}}{d} \overset{\bar{11}}{|} \overset{\bar{8}}{a} \overset{\bar{7}}{|} \overset{\bar{6}}{b} \overset{\bar{4}}{|} \overset{\bar{3}}{d\bar{d}} \overset{\bar{1}}{|} \overset{0}{\bar{a}} \overset{1}{|} .$$

Now all maximal free intervals have length one.

In the next step we show that we can reduce the alphabet of constants to be Γ . The inclusion of Γ into Γ'_0 defines an admissible base change $\beta : \Gamma \rightarrow \Gamma'_0$. Consider $E'_0 = (\Gamma, h, \Omega_0, \rho_0; L_0 = R_0)$ where h is the restriction of the mapping h'_0 . Then we have $\pi^*(E_0) = \beta_*(E'_0)$. The search graph contains an arc from E_0 to E'_0 , since we may choose δ to be the identity. The equation with constraints E'_0 has a solution σ' with $\sigma'(L_0) = w'_0$ and $\exp(w'_0) \leq \exp(w_0)$.

In order to avoid an excess of notation we identify E_0 and E'_0 , hence we also assume $\sigma = \sigma'$ and $w_0 = w'_0$. However, as a reminder that we have changed the alphabet of constants (recall that some words became letters), we prefer to use the symbol Γ rather than Γ_0 . Thus, in what follows we use the following.

Assumption 44 *The input equation E_0 satisfies the following conditions:*

$$\begin{aligned} E_0 &= (\Gamma, h, \Omega_0, \rho_0; L_0 = R_0), \\ L_0 &= x_1 \cdots x_g \text{ and } g \geq 2, \\ R_0 &= x_{g+1} \cdots x_d \text{ and } d > g, \\ |\Gamma| &\leq 2d - 2, \\ |\Omega_0| &\leq 2d. \end{aligned}$$

Moreover: All variables $X \in \Omega_0$ occur in $L_0 R_0 \overline{L_0 R_0}$. There is a solution σ and a word w_0 with $|w_0| = m_0$ and $\exp(w_0) \in 2^{\mathcal{O}(d+n \log n)}$ such that $w_0 = \sigma(L_0) = \sigma(R_0)$ with $\sigma(X_i) \neq 1$ for $1 \leq i \leq d$ and $\rho_0 = h\sigma : \Omega_0 \rightarrow M_{2n} \subseteq \mathbb{B}^{2n \times 2n}$. All maximal free intervals have length exactly one, i.e., every positive interval $[\alpha, \beta]$ with $\beta - \alpha > 1$ contains an implicit cut.

4.10 Critical Words

For each $1 \leq \ell \leq m_0$ we define the set of *critical words* C_ℓ by

$$C_\ell = \{ w_0[\gamma - \ell, \gamma + \ell], w_0[\gamma + \ell, \gamma - \ell] \mid \ell \leq \gamma \leq m_0 - \ell \text{ and } \gamma \text{ is a cut} \}$$

We have $1 \leq |C_\ell| \leq 2d - 4$ and C_ℓ is closed under involution. Each word $u \in C_\ell$ has length 2ℓ , it can be written in the form $u = u_1 u_2$ with $|u_1| = |u_2| = \ell$. The word u_1 (resp. $\overline{u_2}$) appears as a suffix, to the left of some cut and u_2 (resp. $\overline{u_1}$) appears as a prefix, to the right of the same cut.

By B_ℓ we denote the set of triples $(u, w, v) \in (\{1\} \cup \Gamma^\ell) \times \Gamma^+ \times (\{1\} \cup \Gamma^\ell)$ which satisfy the following four conditions:

- 1.) No factor of the word w belongs to C_ℓ .
- 2.) If a factor of the word uwv belongs to C_ℓ , then this factor is a prefix or a suffix of uwv .
- 3.) If $u \neq 1$, then a prefix of uwv of length 2ℓ belongs to C_ℓ ,
- 4.) If $v \neq 1$, then a suffix of uwv of length 2ℓ belongs to C_ℓ .

The set B_ℓ is viewed as a (possibly infinite) alphabet where the involution is defined by $\overline{(u, w, v)} = (\overline{v}, \overline{w}, \overline{u})$. We can define a morphism $\pi_\ell : B_\ell^* \rightarrow \Gamma^*$ by $\pi_\ell(u, w, v) = w \in \Gamma^+$. It is extended to a projection $\pi_\ell : (B_\ell \cup \Gamma)^* \rightarrow \Gamma^*$ by leaving Γ invariant. We define $h_\ell : (B_\ell \cup \Gamma)^* \rightarrow M_{2n}$ by $h_\ell = h\pi_\ell$, i.e., $h_\ell(a) = h(a)$ for $a \in \Gamma$ and $h_\ell(u, w, v) = h(w)$ for $(u, w, v) \in B_\ell$. The symbols π_ℓ and h_ℓ are also used for restrictions of the morphisms π_ℓ and h_ℓ .

Later we consider finite sets $\Gamma_\ell, \Gamma_{\ell, \ell'}$ such that $\Gamma \subseteq \Gamma_\ell \subseteq \Gamma_{\ell, \ell'} \subseteq B_\ell \cup \Gamma$. Then $\pi_{\ell, \ell'} : \Gamma_{\ell, \ell'}^* \rightarrow \Gamma_\ell^*$ denotes the projection given by $\pi_{\ell, \ell'}(u, w, v) = w \in \Gamma^*$ for $(u, w, v) \in \Gamma_{\ell, \ell'} \setminus \Gamma_\ell$ and $\pi_{\ell, \ell'}(u, w, v) = (u, w, v)$ for $(u, w, v) \in \Gamma_\ell$. By $h_\ell : \Gamma_\ell^* \rightarrow M_{2n}$ and $h_{\ell, \ell'} : \Gamma_{\ell, \ell'}^* \rightarrow M_{2n}$ we denote the restrictions of $h_\ell : (B_\ell \cup \Gamma)^* \rightarrow M_{2n}$. We have $h_{\ell, \ell'} = h_\ell \pi_{\ell, \ell'}$.

For every non-empty word $w \in \Gamma^+$ we define its ℓ -factorization as follows. We write

$$F_\ell(w) = (u_1, w_1, v_1) \cdots (u_k, w_k, v_k) \in B_\ell^+$$

such that $w = w_1 \cdots w_k$ and for $1 \leq i \leq k$ the following conditions are satisfied:

- u_i is a suffix of $w_1 \cdots w_{i-1}$,
- $u_i = 1$ if and only if $i = 1$,
- v_i is a prefix of $w_{i+1} \cdots w_k$,

- $v_i = 1$ if and only if $i = k$.

w_1		v_1		$w_2 \cdots w_{k-1}$				u_k		w_k	
				\cdots							
u_2		w_2	v_2	\cdots			u_{k-1}	w_{k-1}	v_{k-1}		

Note that the ℓ -factorization of a word w is unique. For $k \geq 2$ we have $|w_1| \geq \ell$ and $|w_k| \geq \ell$, but all other w_i may be short. If no critical word appears as a factor of w , then $F_\ell(w) = (1, w, 1)$. In particular, this is the case for $|w| < 2\ell$. If we have $w = puvq$ with $|u| = |v| = \ell$ and $uv \in C_\ell$, then there is a unique $i \in \{1, \dots, k-1\}$ such that $u = u_{i+1}$, $v = v_i$, and $pu = w_1 \cdots w_i$, $vq = w_{i+1} \cdots w_k$. Thus, $F_\ell(w)$ contains a factor $(u_i, w_i, v)(u, w_{i+1}, v_{i+1})$ where v is a prefix of $w_{i+1}v_{i+1}$ and u is a suffix of u_iw_i . For example, the ℓ -factorization of $uv \in C_\ell$ with $|u| = |v| = \ell$ is

$$F_\ell(uv) = (1, u, v)(u, v, 1).$$

We define the *head*, *body*, and *tail* of a word w based on its ℓ -factorization

$$F_\ell(w) = (u_1, w_1, v_1) \cdots (u_k, w_k, v_k)$$

in B_ℓ^* and Γ^* as follows:

$$\begin{aligned} \text{Head}_\ell(w) &= (u_1, w_1, v_1) \in B_\ell, \\ \text{head}_\ell(w) &= w_1 \in \Gamma^+, \\ \text{Body}_\ell(w) &= (u_2, w_2, v_2) \cdots (u_{k-1}, w_{k-1}, v_{k-1}) \in B_\ell^*, \\ \text{body}_\ell(w) &= w_2 \cdots w_{k-1} \in \Gamma^*, \\ \text{Tail}_\ell(w) &= (u_k, w_k, v_k) \in B_\ell, \\ \text{tail}_\ell(w) &= w_k \in \Gamma^+. \end{aligned}$$

For $k \geq 2$ (in particular, if $\text{body}_\ell(w) \neq 1$) we have

$$\begin{aligned} F_\ell(w) &= \text{Head}_\ell(w)\text{Body}_\ell(w)\text{Tail}_\ell(w), \\ w &= \text{head}_\ell(w)\text{body}_\ell(w)\text{tail}_\ell(w). \end{aligned}$$

Moreover, u_2 is a suffix of w_1 and v_{k-1} is a prefix of w_k . Assume $\text{body}_\ell(w) \neq 1$ and let $u, v \in \Gamma^*$ be any words. Then we can view w in the context $u w v$ and $\text{Body}_\ell(w)$ appears as a proper factor in the ℓ -factorization of $u w v$. More precisely, let

$$F_\ell(u w v) = (u_1, w_1, v_1) \cdots (u_k, w_k, v_k).$$

Then there are unique $1 \leq p < q \leq k$ such that:

$$\begin{aligned} F_\ell(uwv) &= (u_1, w_1, v_1) \cdots (u_p, w_p, v_p) \text{Body}_\ell(w)(u_q, w_q, v_q) \cdots (u_k, w_k, v_k), \\ w_1 \cdots w_p &= u \text{head}_\ell(w), \\ w_q \cdots w_k &= \text{tail}_\ell(w)v \end{aligned}$$

Finally, we note that the above definitions are compatible with the involution. We have $F_\ell(\bar{w}) = \overline{F_\ell(w)}$, $\text{Head}_\ell(\bar{w}) = \overline{\text{Tail}_\ell(w)}$, and $\text{Body}_\ell(\bar{w}) = \overline{\text{Body}_\ell(w)}$.

4.11 The ℓ -Transformation

By Assumption 44 we have $E_0 = (\Gamma, h, \Omega_0, \rho_0; x_1 \cdots x_g = x_{g+1} \cdots x_d)$. and the equation has a solution σ where $w_0 = \sigma(x_1 \cdots x_g) = \sigma(x_{g+1} \cdots x_d)$ and $m_0 = |w_0|$. We let $1 \leq \ell \leq m_0$ and we consider the ℓ -factorization of the word w_0 :

$$F_\ell(w_0) = (u_1, w_1, v_1) \cdots (u_k, w_k, v_k).$$

A sequence $S = (u_p, w_p, v_p) \cdots (u_q, w_q, v_q)$ with $1 \leq p \leq q \leq k$ is called an ℓ -factor. We say that S is a *cover* of a positive interval $[\alpha, \beta]$, if both, $|w_1 \cdots w_{p-1}| \leq \alpha$ and $|w_{q+1} \cdots w_k| \leq m_0 - \beta$. That is, $w_0[\alpha, \beta]$ is a factor of $w_p \cdots w_q$. It is a *minimal cover*, if neither $(u_{p+1}, w_{p+1}, v_{p+1}) \cdots (u_q, w_q, v_q)$ nor $(u_p, w_p, v_p) \cdots (u_{q-1}, w_{q-1}, v_{q-1})$ is a cover of $[\alpha, \beta]$. The minimal cover exists and it is unique.

We let $\Omega_\ell = \{X \in \Omega_0 \mid \text{body}_\ell(\sigma(X)) \neq 1\}$, and we are going to define a new left-hand side $L_\ell \in (B_\ell \cup \Omega_\ell)^*$ and a new right-hand side $R_\ell \in (B_\ell \cup \Omega_\ell)^*$. For L_ℓ we consider those $1 \leq i \leq g$ where $\text{body}_\ell(\sigma(x_i)) \neq 1$. Note that this implies $x_i \in \Omega_\ell$ since $\ell \geq 1$ and the body of a constant is always empty. Recall the definition of $l(i)$ and $r(i)$, and define $\alpha = l(i) + |\text{head}_\ell(\sigma(x_i))|$ and $\beta = r(i) - |\text{tail}_\ell(\sigma(x_i))|$. We have $w_0[\alpha, \beta] = \text{body}_\ell(\sigma(x_i))$. Next consider the ℓ -factor $S_i = (u_p, w_p, v_p) \cdots (u_q, w_q, v_q)$ which is the minimal cover of $[\alpha, \beta]$. Then we have $1 < p \leq q < k$ and $w_p \cdots w_q = w_0[\alpha, \beta] = \text{body}_\ell(\sigma(x_i))$. The value of S_i depends only on x_i , but not on the choice of the index i . This means $S_i = S_j$ whenever $x_i = x_j$.

We replace the ℓ -factor S_i in $F_\ell(w_0)$ by the variable x_i . Having done this for all $1 \leq i \leq g$ with $\text{body}_\ell(\sigma(x_i)) \neq 1$ we obtain the left-hand side $L_\ell \in (B_\ell \cup \Omega_\ell)^*$ of the ℓ -transformation E_ℓ . For R_ℓ we proceed analogously by replacing those ℓ -factors S_i where $\text{body}_\ell(\sigma(x_i)) \neq 1$ and $g+1 \leq i \leq d$.

For E_ℓ we cannot use the alphabet B_ℓ , because it might be too large (even infinite). Therefore we let Γ'_ℓ be the smallest subset of B_ℓ which is closed under involution and which satisfies $L_\ell R_\ell \in (\Gamma'_\ell \cup \Omega_\ell)^*$.

We let $\Gamma_\ell = \Gamma'_\ell \cup \Gamma$. (We allow Γ because the constants of Γ make it easy to cope with the constraints.) Recall that $h_\ell(u, w, v) = h(w)$ for $(u, w, v) \in \Gamma_\ell \setminus \Gamma$

and $h_\ell(a) = h(a)$ for $a \in \Gamma$. Finally, we define the mapping $\rho_\ell : \Omega_\ell \rightarrow M_{2n}$ by $\rho_\ell(X) = h(\text{body}_\ell(\sigma(X)))$. The reason is that we know $\rho(X) = h(\sigma(X))$. We can write $\sigma(X) = u\text{body}_\ell(\sigma(X))v$, hence $h(\sigma(X)) = h(u)\rho_\ell(X)h(v)$. The steps above define the ℓ -transformation and yield the following equation:

$$E_\ell = (\Gamma_\ell, h_\ell, \Omega_\ell, \rho_\ell; L_\ell = R_\ell).$$

Example 45 We continue with our example $aX\bar{X}\bar{a} = Y\bar{b}Y\bar{a}b\bar{Y}$ and the solution σ which has been given by

$$w_0 = | a | d\bar{d} | \bar{b} | ad | \bar{d} | \bar{a} | b | d\bar{d} | \bar{a} |,$$

where the bars show the cuts.

Up to involution, the set C_1 is given by $\{ad, bd, \bar{a}\bar{b}, d\bar{d}\}$ and C_2 is given by $\{d\bar{d}\bar{b}a, \bar{d}\bar{b}ad, add\bar{a}, d\bar{d}\bar{a}b\}$. The 1-factorization of w_0 can be obtained letter by letter.

The 2-factorization of w_0 is given by the following sequence:

$$(1, add, \bar{b}a)(d\bar{d}, \bar{b}, ad)(\bar{d}\bar{b}, ad, \bar{d}\bar{a})(ad, \bar{d}, \bar{a}b)(d\bar{d}, \bar{a}, bd)(\bar{d}\bar{a}, b, d\bar{d})(\bar{a}b, d\bar{d}\bar{a}, 1).$$

Recall $\sigma(X) = d\bar{d}\bar{b}ad$ and $\sigma(Y) = add\bar{d}$. Hence their 2-factorizations are $(1, d\bar{d}, \bar{b}a)(d\bar{d}, \bar{b}, ad)(\bar{d}\bar{b}, ad, 1)$ and $(1, add, 1)$, respectively.

Let us rename the letters:

$$\begin{aligned} a &= (1, add, \bar{b}a) \\ b &= (\bar{d}\bar{a}, b, d\bar{d}) \\ c &= (\bar{d}\bar{b}, ad, \bar{d}\bar{a}) \\ d &= (ad, \bar{d}, \bar{a}b) \\ e &= (d\bar{d}, \bar{a}, bd) \end{aligned}$$

After this renaming the 2-factorization of w_0 becomes $\bar{a}b\bar{c}d\bar{e}b\bar{a}$ and the equation E reduces to $E_2 : aXcde\bar{X}\bar{a} = \bar{a}b\bar{c}d\bar{e}b\bar{a}$ since the body of $\sigma(Y)$ is empty. The reader can check that the 3-factorization of w_0 after renaming is the very same word as the 2-factorization, but the 3-factorization of $\sigma(X)$ is now one letter, $(1, d\bar{d}\bar{b}ad, 1)$, so E_3 becomes a trivial equation. Plandowski's algorithm will return true at this stage.

Remark 46 i) In the extreme case $\ell = m_0$, the ℓ -transformation becomes trivial. Let $a = (1, w_0, 1)$. Then $\bar{a} = (1, \bar{w}_0, 1)$ and $\Gamma_{m_0} = \{a, \bar{a}\} \cup \Gamma$. Moreover, we have $L_{m_0} = R_{m_0} = a$, and $h_{m_0}(a) = h(w_0) \in M_{2n}$. Since $\Omega_{m_0} = \emptyset$, the equation with constraints E_{m_0} trivially has a solution. It is clear that E_{m_0}

is a node in the search graph, and if we reach E_{m_0} , then the algorithm will return true.

ii) The other extreme case is $\ell = 1$. We develop the technical details as an example. Consider a triple $(u, w, v) \in \Gamma_1$ which appears in $F_1(w_0)$. Then $w = w_0[\alpha, \beta]$ for some $\beta - \alpha \geq 1$. All maximal free intervals have length 1 (by Assumption 44). Assume $\beta - \alpha \geq 2$, then $[\alpha, \beta]$ would contain an implicit cut γ and $w_0[\gamma - 1, \gamma + 1] \in C_1$. But no critical word is a factor of w , $\beta - \alpha = 1$. An immediate consequence is $|\Gamma_1| \leq (|\Gamma| + 1)^3 \in \mathcal{O}(d^3)$, since $|\Gamma| \leq 2d - 2$. (More precisely, we could bound $|\Gamma_1|$ by $6d$, but $|\Gamma_1| \in \mathcal{O}(d^3)$ is good enough for our purpose.) Let $X \in \Omega_0$. Then $\text{Body}_1(\sigma(X)) \neq 1$ if and only if $|\sigma(X)| \geq 3$. Thus, for $X \in \Omega_1$ we have $\sigma(X) = bcu = vde$ with $b, c, d, e \in \Gamma$ and $u, v \in \Gamma^+$. It follows:

$$F_1(\sigma(X)) = (1, b, c)(b, c, v_2) \cdots (u_{|v|+1}, d, e)(d, e, 1).$$

For example, for $|v| = 1$ this means $b = u_{|v|+1}$, $c = d$, and $v_2 = e$.

We can describe $L_1 \in \Gamma_1^*$ as follows:

For $1 \leq i \leq g$ let $w_i = \sigma(x_i)$ and a_i the last letter of $\sigma(x_{i-1})$ if $i > 1$ and $a_1 = 1$. Let f_i the first letter of $\sigma(x_{i+1})$ if $i < g$ and $f_g = 1$. Let b_i the first letter of w_i and e_i the last letter of w_i .

For $|w_i| = 1$ we replace x_i by the 1-factor (a_i, b_i, f_i) .

For $|w_i| = 2$ we replace x_i by the 1-factor $(a_i, b_i, e_i)(b_i, e_i, f_i)$.

For $|w_i| \geq 3$ we let c_i be the second letter of w_i and d_i its second last. In this case we replace x_i by $(a_i, b_i, c_i)x_i(d_i, e_i, f_i)$.

The definition of R_1 is analogous. Thus, we obtain $|L_1 R_1| \leq 3|L_0 R_0| = 3d$, and E_1 is admissible.

By Remark 46 the equations E_1 and E_{m_0} are admissible and hence nodes of the search graph of E_0 . The goal is to reach E_{m_0} , but it is not clear yet, neither that the ℓ -transformations with $1 < \ell < m_0$ belong to the search graph nor that there are arcs from E_0 to E_1 or from E_1 to E_2 and so on. We prove these statements in the next sections.

4.12 The ℓ -transformation E_ℓ is Admissible

Proposition 47 *There is a polynomial p_0 (of degree 4) such that each E_ℓ is admissible with respect to p_0 for all $\ell \geq 1$.*

Proof. The input size is $d+n+\log_2(|\Gamma|+|\Omega_0|)$. We have $|\Gamma|+|\Omega_0| \leq 4d-2$ and $E_0 = (\Gamma, h, \Omega_0, \rho_0; x_1 \cdots x_g = x_{g+1} \cdots x_d)$. The constraints are Boolean $n \times n$ -matrices and d is the length of the equation. It is enough to show that L_ℓ and R_ℓ can be represented by exponential expressions of size $\mathcal{O}(d^2(d+n \log n))$.

Then Γ_ℓ can have size at most $\mathcal{O}(d^2(d + n \log n))$ and the assertion follows. We will estimate the size of an exponential expression for L_ℓ , only.

We start again with the ℓ -transformation

$$F_\ell(w_0) = (u_1, w_1, v_1) \cdots (u_k, w_k, v_k).$$

If k is small there is nothing to do since $|L_\ell| \leq |F_\ell(w_0)|$. An easy reflection shows that $|L_\ell|$ can become large, only if there is some $1 \leq i \leq g$ such that $\text{head}_\ell(\sigma(x_i))$ or $\text{tail}_\ell(\sigma(x_i))$ is long. By symmetry we treat the case $\text{head}_\ell(\sigma(x_i))$ only and we fix some notation. We let $1 \leq i \leq g$, $\alpha = l(i)$, and $\beta = \alpha + |\text{head}_\ell(\sigma(x_i))|$. Let

$$(u_{p-1}, w_{p-1}, v_{p-1}) \cdots (u_{q+1}, w_{q+1}, v_{q+1})$$

be a minimal cover of $[\alpha, \beta]$. (The definition of a minimal cover has been given at the beginning of Subsection 4.11.) We may assume that $q - p$ is large. It is enough to show that the ℓ -factor

$$(u_p, w_p, v_p) \cdots (u_q, w_q, v_q)$$

has an exponential expression of size in $\mathcal{O}(d(d + n \log n))$, because we want the whole expression to have size in $\mathcal{O}(d^2(d + n \log n))$.

Note that $w_p \cdots w_q$ is a proper factor of $\text{head}_\ell(\sigma(x_i))$. Hence no critical word of C_ℓ can appear as a factor inside $w_p \cdots w_q$. This means there is some $p \leq s \leq q$ such that both, $|w_p \cdots w_{s-1}| < \ell$ and $|w_{s+1} \cdots w_q| < \ell$. Indeed, if $|w_p \cdots w_{q-1}| < \ell$, then we choose $s = q$. Otherwise we let $p \leq s \leq q$ be minimal such that $|w_p \cdots w_s| \geq \ell$. Then $|w_{s+1} \cdots w_q| \geq \ell$ is impossible because $u_{s+1}v_s \in C_\ell$ would appear as a factor in $w_p \cdots w_q$. We can write

$$(u_p, w_p, v_p) \cdots (u_q, w_q, v_q) = S_1(u_s, w_s, v_s)S_2.$$

Since $(u_s, w_s, v_s) \in \Gamma_\ell$ is a letter, it is enough to show that there are exponential expressions for S_i of size $\mathcal{O}(d(d + n \log n))$ for $i = 1, 2$. This follows from Lemma 48 with $c = 1$. \square

The statement of Lemma 48 is more general than needed for the proof of Proposition 47, but later it is used for other values of c . In fact, it will be used for $c \leq 32d$.

Lemma 48 *Let $c > 0$ be a value which might depend on d (and n) and let*

$$S = (u_1, w_1, v_1) \cdots (u_k, w_k, v_k) \in B_\ell^*$$

be a sequence which appears as some ℓ -factor in $F_\ell(w_0)$. If we have $k \leq 3$ or $|w_2 \cdots w_{k-1}| \leq c\ell$, then the sequence S can be represented by some exponential expression of size $\mathcal{O}(cd(d + n \log n))$.

Proof. Clearly, we may assume $k > 3$. We show that there is an exponential expression of size $\mathcal{O}(d(d + n \log n))$ under the assumption $|w_1 \cdots w_k| < \ell$. (Note that c has been removed from the \mathcal{O} -term.) This is enough, because we can write S as $a_0 S_1 a_1 \cdots S_{c'} a_{c'}$, where $c' \leq c$, the a_i are letters, and each S_i satisfies the assumption. Due to the factorization we may also assume $u_1 \neq 1 \neq v_k$ and therefore we may define u_{k+1} as the suffix of length ℓ of $u_1 w_1 \cdots w_k$. For $1 \leq i \leq k$ let $z_i = u_{i+1} v_i$. Then $z_i \in C_\ell$ is a critical word which appears as a factor in $z = u_1 w_1 w_2 \cdots w_k v_k$. If the words z_i , $1 \leq i < k$ are pairwise different, then $k - 1 \leq |C_\ell| \in \mathcal{O}(d)$ and we are done. Hence we may assume that there are repetitions. Let j be the smallest index such that a critical word is seen for the second time and let $i < j$ be the first appearance of z_j . This means for $1 \leq i < j$ the words z_1, \dots, z_{j-1} are pairwise different and $z_i = z_j$. Now, $|w_1 \cdots w_k| < \ell$ and $|z_i| = 2\ell$, hence z_i and z_j overlap in z . We can choose r maximal such that $u_1 w_1 \cdots w_i (w_{i+1} \cdots w_j)^r v_j$ is a prefix of the word z . (Note that the last factor v_j insures that the prefix ends with z_j .) For some index $s > j$ we can write

$$z = u_1 w_1 \cdots w_i (w_{i+1} \cdots w_j)^r w_s \cdots w_k v_k.$$

We claim that $z_i \notin \{z_s, \dots, z_k\}$. Indeed, let t be maximal such that $z_i = z_t$ and assume that $j \neq t$. Then both, $|w_{i+1} \cdots w_j|$ and $|w_{j+1} \cdots w_t|$ are periods of z_i , but $|w_{i+1} \cdots w_t| \leq |z_i|$. Hence by Fine and Wilf's Theorem [23] we obtain that the greatest common divisor of $|w_{i+1} \cdots w_j|$ and $|w_{j+1} \cdots w_t|$ is a period, too. Due to the definition of an ℓ -factorization (z_j was the first repetition) the length $|w_{j+1} \cdots w_t|$ is therefore a multiple of $|w_{i+1} \cdots w_j|$ and we must have $t = s - 1$. This shows the claim. Moreover, we have

$$\begin{aligned} & (u_1, w_1, v_1) \cdots (u_k, w_k, v_k) \\ &= (u_1, w_1, v_1) \cdots (u_i, w_i, v_i) [(u_{i+1}, w_{i+1}, v_{i+1}) \cdots (u_j, w_j, v_j)]^r S' \end{aligned}$$

where $S' = (u_s, w_s, v_s) \cdots (u_k, w_k, v_k)$ for $s = i + 1 + r(j - i)$ and $r \geq 1$. We have $r \leq \exp(w_0)$, hence $r \in 2^{\mathcal{O}(d+n \log n)}$. It follows that

$$(u_1, w_1, v_1) \cdots (u_i, w_i, v_i) [(u_{i+1}, w_{i+1}, v_{i+1}) \cdots (u_j, w_j, v_j)]^r$$

is an exponential expression of size $j + \lceil \log_2(r) \rceil \in \mathcal{O}(d + n \log n)$. More precisely, we can effectively calculate some constant \tilde{c} such that $j + \lceil \log_2(r) \rceil \leq \tilde{c}(d + n \log_2 n)$.

We have $|\{z_s, \dots, z_k\}| < |\{z_1, \dots, z_k\}|$. Therefore by induction we may assume that the sequence $S' = (u_s, w_s, v_s) \cdots (u_k, w_k, v_k)$ has an exponential expression of size at most $|\{z_s, \dots, z_k\}| \tilde{c}(d + n) \log_2 n$. Hence the exponential expression for S has size at most

$$\tilde{c}(d + n \log_2 n) + |\{z_s, \dots, z_k\}| \tilde{c}(d + n \log_2 n) \leq |\{z_1, \dots, z_k\}| \tilde{c}(d + n \log_2 n).$$

Thus, the size is in $\mathcal{O}(d(d + n \log n))$. □

Remark 49 *At this stage we know that all ℓ -transformations are admissible with respect to some suitable polynomial p_0 of degree 4. Next we show that we can modify the polynomial p_0 such that the search graph also contains arcs $E_0 \rightarrow E_1$ and $E_\ell \rightarrow E_{\ell'}$ for $1 \leq \ell < \ell' \leq 2\ell$. For this reason we use the notion of admissibility with respect to the 4-th power p_0^4 of p_0 . Thus, admissibility is meant with respect to a polynomial of degree 16.*

4.13 The Arc from E_0 to E_1

We present the formal construction of the arc from E_0 to E_1 . We give all technical details since this arc is the model for the more complicated way the other arcs are constructed in the search graph.

An explicit description of $E_1 = (\Gamma_1, h_1, \Omega_1, \rho_1; L_1 = R_1)$ has been given in Remark 46. The letters of Γ_1 can be written either as (a, b, c) or as b with $a, c \in \Gamma \cup \{1\}$ and $b \in \Gamma$. We define an admissible base change $\beta : \Gamma_1 \rightarrow \Gamma$ by $\beta(a, b, c) = b$ and $\beta(b) = b$ for $b \in \Gamma$. Trivially, $h_1 = h\beta$. Define $E_{0,1} = \beta_*(E_1)$. Then we have $L_{0,1} = \beta(L_1)$ and $R_{0,1} = \beta(R_1)$ where $\beta : (\Gamma_1 \cup \Omega_1)^* \rightarrow (\Gamma \cup \Omega_1)^*$ is the extension with $\beta(X) = X$ for all $X \in \Omega_1$. We have $\Gamma_{0,1} = \Gamma$.

It is now obvious how to define the partial solution $\delta : \Omega_0 \rightarrow \Gamma\Omega_1\Gamma \cup \Gamma^*$ such that $\delta_*(E_0) = E_{0,1}$. If $|\sigma(X)| \leq 2$, then we let $\delta(X) = \sigma(X)$. For $|\sigma(X)| \geq 3$ we write $\sigma(X) = aub$ with $a, b \in \Gamma$ and $u \in \Gamma^+$. Then we have $X \in \Omega_1 = \Omega_{0,1}$ and we define $\delta(X) = aXb$ and $\rho_{0,1}(X) = h(u)$. For $X \in \Omega_1$ we have, by definition, $\rho_1(X) = h(\text{body}_1(\sigma(X)))$, hence $\rho_{0,1} = \rho_1$, too. This shows that, indeed, $\delta_*(E_0) = \beta_*(E_1)$. Formally, we can write this as $\delta_*(\pi^*(E_0)) = \beta_*(E_1)$, where π is the identity. This yields the arc from E_0 to E_1 .

4.14 The Equation $E_{\ell,\ell'}$ for $1 \leq \ell < \ell' \leq 2\ell$

In order to establish the arcs from E_ℓ to $E_{\ell'}$ for all $1 \leq \ell < \ell' \leq 2\ell$ we use an intermediate equation $E_{\ell,\ell'}$ such that there is an admissible base change β , a projection π , and a partial solution δ with

$$\delta_*(\pi^*(E_\ell)) \equiv E_{\ell,\ell'} = \beta_*(E_{\ell'}).$$

The way we move from E_ℓ to $E_{\ell'}$ is visualized by Figure 1.

We begin with the definition of the base change β . Recall $\Gamma \subseteq \Gamma_{\ell'} \subseteq B_{\ell'} \cup \Gamma$. As expected, we define $\beta(a) = a$ for $a \in \Gamma$. Consider some $(u, w, v) \in \Gamma_{\ell'} \setminus \Gamma$.




Figure 1: The search graph and its neighborhood


It is enough to define $\beta(u, w, v)$ or $\beta(\bar{v}, \bar{w}, \bar{u})$. Hence we may assume that (u, w, v) appears in the ℓ' -factorization $F_{\ell'}(w_0)$. Therefore we find a positive interval $[\alpha_0, \beta_0]$ such that $w = w_0[\alpha_0, \beta_0]$ and such that the following two conditions are satisfied:

- 1.) We have $u = 1$ and $\alpha_0 = 0$ or $|u| = \ell'$, $\alpha_0 \geq \ell'$, and $u = w_0[\alpha_0 - \ell', \alpha_0]$.
- 2.) We have $v = 1$ and $\beta_0 = m_0$ or $|v| = \ell'$, $\beta_0 \leq m_0 - \ell'$, and $v = w_0[\beta_0, \beta_0 + \ell']$.

Let $(u_p, w_p, v_p) \cdots (u_q, w_q, v_q)$ be the ℓ -factor which is the minimal cover of $[\alpha_0, \beta_0]$ with respect to the ℓ -factorization $F_{\ell}(w_0)$. Since $\ell \leq \ell'$ we have $w_p \cdots w_q = w$. Moreover, the word u_p is a suffix of u and v_q is a prefix of v . We define

$$\beta(u, w, v) = (u_p, w_p, v_p) \cdots (u_q, w_q, v_q) \in B_{\ell}^+.$$

We have the following picture:



The definition does not depend on the choice of $[\alpha_0, \beta_0]$ as long as $0 \leq \alpha_0 < \beta_0 \leq m_0$ and 1.) and 2.) are satisfied. We have $\beta(u, w, v) = \beta(\bar{v}, \bar{w}, \bar{u})$

and $h_\ell\beta = h_{\ell'}$. Now let $\Gamma_{\ell,\ell'} \subseteq B_\ell \cup \Gamma$ be the smallest subset such that $\beta(\Gamma_{\ell'}) \subseteq \Gamma_{\ell,\ell'}^*$. Then $\Gamma_{\ell,\ell'}$ contains Γ and it is closed under involution (since $\Gamma_{\ell'}$ has this property). An easy reflection shows that $\Gamma_\ell \subseteq \Gamma_{\ell,\ell'}$. This will become essential in Subsection 4.15.

We view β as a morphism $\beta : \Gamma_{\ell'}^* \rightarrow \Gamma_{\ell,\ell'}^*$ and we have $h_{\ell,\ell'}\beta = h_\ell$. Define $E_{\ell,\ell'} = \beta_*(E_{\ell'})$. Then

$$E_{\ell,\ell'} = (\Gamma_{\ell,\ell'}, h_{\ell,\ell'}, \Omega_{\ell'}, \rho_{\ell'}; \beta(L_{\ell'}) = \beta(R_{\ell'}).$$

Let us show that β is admissible. Since $E_{\ell'}$ is already known to be admissible with respect to some polynomial of degree 4, it is enough to find some admissible exponential expression (again with respect to some polynomial of degree 4) for the ℓ -factor above

$$\beta(u, w, v) = (u_p, w_p, v_p) \cdots (u_q, w_q, v_q).$$

Using the same terminology as above there is some positive interval $[\alpha_0, \beta_0]$ such that $w_p \cdots w_q = w_0[\alpha_0, \beta_0]$, the word u is a suffix of $w_0[0, \alpha_0]$, and v is a prefix of $w_0[\beta_0, m_0]$. If $q - p$ is small, there is nothing to do. By Lemma 48 we may also assume that $\beta_0 - \alpha_0 > 32d\ell$. We inductively define a sequence of positions

$$\alpha_0 \leq \alpha_1 \leq \cdots \leq \alpha_i \leq \cdots \leq \beta_i \leq \cdots \leq \beta_1 \leq \beta_0.$$

In each step we let $W_i = w_0[\alpha_i, \beta_i]$. Thus, $W_0 = w_p \cdots w_q$. Assume that $W_i = w_0[\alpha_i, \beta_i]$ is already defined such that $\beta_i - \alpha_i \geq 2$. The interval $[\alpha_i, \beta_i]$ is not free. Hence, there is some implicit cut γ_i with $\alpha_i < \gamma_i < \beta_i$. The word W_i is a factor of w , hence no factor of W_i belongs to the set of critical words $C_{\ell'}$. This implies $\beta_i - \gamma_i < \ell'$ or $\gamma_i - \alpha_i < \ell'$. If we have $\beta_i - \gamma_i < \ell'$ then we let $\alpha_{i+1} = \alpha_i$ and $\beta_{i+1} = \gamma_i$. In the other case we let $\alpha_{i+1} = \gamma_i$ and $\beta_{i+1} = \beta_i$. Thus W_{i+1} is defined such that W_{i+1} is a proper factor of W_i with $|W_i| - |W_{i+1}| < \ell'$.

We need some additional book keeping. We define $r_i \in \{1, r\}$ by $r_i = r$ if $\beta_i = \beta_{i+1}$ and $r_i = 1$ otherwise (i.e., $\alpha_i = \alpha_{i+1}$). Furthermore the implicit cut γ_i corresponds to some real cut γ'_i and $\alpha'_i < \gamma'_i < \beta'_i$ such that $W_i = w_0[\alpha'_i, \beta'_i]$ or $W_i = w_0[\beta'_i, \alpha'_i]$. We define $s_i \in \{+, -\}$ by $s_i = +$ if $W_i = w_0[\alpha'_i, \beta'_i]$ and $s_i = -$ otherwise (in particular, $s_i = -$ implies $\overline{W_i} = w_0[\alpha'_i, \beta'_i]$). The triple (γ'_i, r_i, s_i) is denoted by $\gamma(i)$. There are at most $4(d-2)$ such triples and $\gamma(i)$ is defined whenever W_{i+1} is defined. We stop the induction procedure after the first repetition of some $\gamma(i)$. Thus we find $0 \leq i < j < 4d$ such that $\gamma(i) = \gamma(j)$. We obtain a sequence $W_0, W_1, \dots, W_i, \dots, W_j$ where each word is a proper factor of the preceding one. We have $|W_0| - |W_j| < 4d\ell' \leq 8d\ell$ and due to $|W_0| > 32d\ell$ the sequence above really exists, moreover $|W_j| > 24d\ell$.

Next, we show that W_j has a non-trivial overlap with itself. We treat the case $\gamma(i) = \gamma(j) = (\gamma, r, +)$ only. The other three cases $(\gamma, r, -)$, $(\gamma, l, +)$, and $(\gamma, l, -)$ can be treated analogously. For some $\alpha' < \gamma < \beta'$ we have $W_i = w_0[\alpha', \beta']$ and $W_{i+1} = w_0[\gamma, \beta']$. Thus, for some $\gamma \leq \mu < \nu \leq \beta'$ we have $W_j = w_0[\mu, \nu]$ and we can assume that $\mu - \gamma < (j - i)\ell' \leq 4d\ell' - \ell' \leq 8d\ell - \ell'$. On the other hand we have $\gamma(j) = (\gamma, r, +)$, too. Hence for some $\mu' < \gamma < \nu'$ with $\gamma - \mu' < \ell'$ we have $W_j = w_0[\mu', \nu']$, too. Therefore $0 < \mu - \mu' < 8d\ell$ and W_j has some non-trivial overlap. We may choose $W = w_0[\mu', \mu]$ and it follows that we can write $W_j = W^e W'$ such that $1 \leq |W| < 8d\ell$ and W' is a prefix of W .

Putting everything together, we arrive in all cases at a factorization $W_0 = UW^eV$ with $e \leq \exp(w_0)$, $1 \leq |W| < 8d\ell$, and $|U| + |V| < 16d\ell$.

We have not finished yet. Recall that we are looking for an admissible exponential expression for

$$\beta(u, w, v) = (u_p, w_p, v_p) \cdots (u_q, w_q, v_q).$$

Due to $|W_0| > \ell$ we can choose r minimal, $p < r \leq q + 1$, and s maximal $p - 1 \leq s < q$ such that $|w_p \cdots w_{r-1}| > |U| + \ell$ and $|w_{s+1} \cdots w_q| > |V| + \ell$. By Lemma 48 we may assume $r < s$ and it is enough to find an exponential expression for

$$S = (u_r, w_r, v_r) \cdots (u_s, w_s, v_s).$$

Note that the word $u_r w_r w_{r+1} \cdots w_s v_s$ is a factor of W^e . Hence we may factorize $W = W'W''$ in such a way that after replacing W by $W''W'$, we may assume that $u_r w_r w_{r+1} \cdots w_s v_s$ is in fact a prefix of W^e . Furthermore, we may assume that $w_r w_{r+1} \cdots w_s > 32d\ell$ and by symmetry we may choose some positive interval $[\alpha_0, \beta_0]$ such that $w_0[\alpha_0, \beta_0] = u_r w_r w_{r+1} \cdots w_s v_s$. Clearly, we have $w_0[i, j] = w_0[i + |W|, j + |W|]$ for all $\alpha_0 \leq i < j \leq \beta_0 - |W|$. In particular, the critical word $w_0[\alpha_0, \alpha_0 + 2\ell]$ appears as $w_0[\alpha_0 + |W|, \alpha_0 + 2\ell + |W|]$ again. This means that there is some $r \leq t < s$ such that $|w_r \cdots w_t| = |W|$. More precisely, we can choose $r \leq t < t' \leq s$ and a maximal $e' \leq e$ such that

$$S = ((u_r, w_r, v_r) \cdots (u_t, w_t, v_t))^{e'} (u_{t'}, w_{t'}, v_{t'}) \cdots (u_s, w_s, v_s).$$

Since $e' \leq \exp(w_0)$, $|w_r \cdots w_t| = |W|$, and $|w_{t'} \cdots w_s| \leq |W|$, the existence of an admissible exponential expression for $\beta(u, w, v)$ follows. Hence β is an admissible base change.

4.15 Passing from E_ℓ to $E_{\ell, \ell'}$ for $1 \leq \ell < \ell' \leq 2\ell$

In the final step we have to show that there exists some projection $\pi : \Gamma_{\ell, \ell'}^* \rightarrow \Gamma_\ell^*$ and some partial solution $\delta : \Omega_\ell \rightarrow \Gamma_{\ell, \ell'}^* \Omega_{\ell'} \Gamma_{\ell, \ell'}^* \cup \Gamma_{\ell, \ell'}^*$ such that

$\delta_*(\pi^*(E_\ell)) \equiv E_{\ell,\ell'}$. We don't have to worry about admissibility anymore. Once $\delta_*(\pi^*(E_\ell)) \equiv E_{\ell,\ell'}$ is established, Theorem 8 is proved.

For the definition of the projection π consider a letter in $\Gamma_{\ell,\ell'} \setminus \Gamma_\ell$. Such a letter has the form $(u, w, v) \in B_\ell$ with $w \in \Gamma^+$. There is no length bound on w known (or needed). We define $\pi(u, w, v) = w$ and this is possible since $\Gamma \subseteq \Gamma_\ell$.

Clearly $\pi(\overline{(u, w, v)}) = \overline{\pi(u, w, v)}$ and $h_{\ell,\ell'}(u, w, v) = h(w) = h_\ell(\pi(u, w, v))$. Thus, $\pi : \Gamma_{\ell,\ell'}^* \rightarrow \Gamma_\ell^*$ defines a projection such that

$$\pi^*(E_\ell) = (\Gamma_{\ell,\ell'}, h_{\ell,\ell'}, \Omega_\ell, \rho_\ell; L_\ell = R_\ell).$$

We have to define a partial solution $\delta : \Omega_\ell \rightarrow \Gamma_{\ell,\ell'}^* \Omega_{\ell'} \Gamma_{\ell,\ell'}^* \cup \Gamma_{\ell,\ell'}^*$ such that $\delta(L_\ell) = \beta(L_{\ell'})$ and $\delta(R_\ell) = \beta(R_{\ell'})$. For this, we have to consider a variable $X \in \Omega$ with $\text{body}_\ell(\sigma(X)) \neq 1$. By symmetry, we may assume that $X = x_i$ for some $1 \leq i \leq g$. Hence $\sigma(X) = w_0[l(i), r(i)]$.

Let $\alpha_X = l(i) + |\text{head}_\ell(\sigma(X))|$ and $\beta_X = r(i) - |\text{tail}_\ell(\sigma(X))|$. Then $l(i) + \ell \leq \alpha_X < \beta_X \leq r(i) - \ell$. Let $(u_p, w_p, v_p) \cdots (u_q, w_q, v_q)$ be the minimal cover of $[\alpha_X, \beta_X]$ with respect to the ℓ -factorization. We have $w_p \cdots w_q = \text{body}_\ell(\sigma(X))$.

For $\text{body}_{\ell'}(\sigma(X)) = 1$ we have $X \in \Omega_\ell \setminus \Omega_{\ell'}$ and we define

$$\delta(X) = (u_p, w_p, v_p) \cdots (u_q, w_q, v_q).$$

Then $\delta(X) \in B_\ell^*$ and $h_\ell \delta(X) = \rho_\ell(X)$ since $\rho_\ell(X) = h(\text{body}_\ell(\sigma(X)))$. It is also clear that the definition does not depend on the choice of i , and we have $\delta(\overline{X}) = \overline{\delta(X)}$.

Recall the definition of $L_{\ell'}$. Since $\text{body}_{\ell'}(\sigma(X)) = 1$, there is a factor $f_1 \cdots f_r$ of $L_{\ell'}$ which belongs to $\Gamma_{\ell'}^*$ and $f_1 \cdots f_r$ covers $[\alpha_X, \beta_X]$ with respect to the ℓ' -factorization $F_{\ell'}(w_0)$. It follows that $\delta(X)$ is a factor of $\beta(f_1 \cdots f_r)$, hence $\delta(X) \in \Gamma_{\ell,\ell'}^*$ by definition of $\Gamma_{\ell,\ell'}$.

For $\text{body}_{\ell'}(\sigma(X)) \neq 1$ we have $X \in \Omega_{\ell'}$ and we find positions $\mu < \nu$ such that $\mu = l(i) + |\text{head}_{\ell'}(\sigma(X))|$ and $\nu = r(i) - |\text{tail}_{\ell'}(\sigma(X))|$.

For some $p \leq r \leq s \leq q$ we have $w_0[\alpha_X, \mu] = w_p \cdots w_{r-1}$, $w_0[\nu, \beta_X] = w_{s+1} \cdots w_q$, and $\text{body}_{\ell'}(\sigma(X)) = w_r \cdots w_s$. We define

$$\delta(X) = (u_p, w_p, v_p) \cdots (u_{r-1}, w_{r-1}, v_{r-1}) X (u_{s+1}, w_{s+1}, v_{s+1}) \cdots (u_q, w_q, v_q).$$

As above, we can verify that $\delta(X) = UXV$ with $U, V \in \Gamma_{\ell,\ell'}^*$ such that $\delta(\overline{X}) = \overline{V X U}$ and $\rho_\ell(X) = h_{\ell,\ell'}(U) \rho_{\ell'}(X) h_{\ell,\ell'}(V)$. Finally, $\delta(L_\ell) = \beta(L_{\ell'})$ and $\delta(R_\ell) = \beta(R_{\ell'})$. Hence $\delta_*(\pi^*(E_\ell)) \equiv \beta_*(E_{\ell'})$. The final step in proving Theorem 8 is completed.

5 Concluding Remarks

The PSPACE-hardness stated in Theorems 3, 5, and 8 is due to rational constraints, but this is a side effect and a nice coincidence, only. The reason to include constraints has been motivated by possible applications to free partially commutative groups (graph groups). When Matiyasevich showed in 1996 that the existential theory of equations in free partially commutative monoids (trace monoids) is decidable [29, 30, 7], it became clear by his method that regular constraints are a powerful tool in order to extend decidability results to other algebraic structures and, in particular, they are necessary for extending Makanin's result from free groups to graph groups. At that time the result of Schulz [38] on word equations with regular constraints had been available but no such analogue for equations in free groups was known. So, the idea was to look for such an analogue first. Inspired by Gutiérrez [15] we were finally led to investigate free monoids with involution and regular constraints. This approach turned out to be fruitful. Based on Theorem 5 of this paper (the results date back to the year 2000) it is shown in [8] that the existential theory of equations in graph groups is decidable. This result is a common generalization of Matiyasevich's decidability result on trace monoids and Makanin's result on free groups. In a continuation of this work on graph groups we obtained various other decidability results about the existential and positive theories in graph products, see [5, 6]. Makanin has also shown that the positive theory in free groups is decidable, [26]. It remains decidable with recognizable constraints [6]. In contrast, the positive theory of equations with rational constraints is undecidable in free groups, because the positive $\forall\exists^3$ -theory of word equations is undecidable [27, 9] and Σ^* is a rational subset of the free group $F(\Sigma)$. So the question remains under which restricted type of constraints the positive theory of equations in free groups remains decidable.

6 Appendix: Proof of Proposition 15

We first repeat the statement of Proposition 15:

Let $E = (\Gamma, h, \Omega, \rho; L = R)$ be a solvable equation with constraints. Then there is a solution $\sigma : \Omega \rightarrow \Gamma^*$ such that $\exp(\sigma(L)) \in 2^{\mathcal{O}(d+n \log n)}$.

Proof. Let $p \in \Gamma^+$ be a primitive word. This means that $p \neq r^k$ for all $k > 1$ and $r \in \Gamma^*$. In the following we also write p^{-1} instead of \bar{p} . Then, p^{-3} for example means the same as \bar{p}^3 . The definition of the p -stable normal form depends on whether or not \bar{p} is a factor of p^2 . So we distinguish two cases.

First case: We assume that \bar{p} is not a factor of p^2 . The idea is to replace each maximal factor of the form p^α with $\alpha \geq 2$ by a sequence $p, \alpha - 2, p$ and each maximal factor of the form \bar{p}^α with $\alpha \geq 2$ by a sequence $\bar{p}, -(\alpha - 2), \bar{p}$. The p -stable normal form (first kind) of $w \in \Gamma^*$ is a shortest sequence (k is minimal)

$$(u_0, \varepsilon_1 \alpha_1, u_1, \dots, \varepsilon_k \alpha_k, u_k)$$

such that $k \geq 0$, $u_0, u_i \in \Gamma^*$, $\varepsilon_i \in \{+1, -1\}$, $\alpha_i \geq 0$ for $1 \leq i \leq k$, and the following conditions are satisfied:

- $w = u_0 p^{\varepsilon_1 \alpha_1} u_1 \cdots p^{\varepsilon_k \alpha_k} u_k$.
- $k = 0$ if and only if neither p^2 nor \bar{p}^2 is a factor of w .
- If $k \geq 1$, then:

$$\begin{aligned} u_0 &\in \Gamma^* p^{\varepsilon_1} \setminus \Gamma^* p^{\pm 2} \Gamma^*, \\ u_i &\in (\Gamma^* p^{\varepsilon_{i+1}} \cap p^{\varepsilon_i} \Gamma^*) \setminus \Gamma^* p^{\pm 2} \Gamma^* \text{ for } 1 \leq i < k, \\ u_k &\in p^{\varepsilon_k} \Gamma^* \setminus \Gamma^* p^{\pm 2} \Gamma^*. \end{aligned}$$

If $(u_0, \varepsilon_1 \alpha_1, u_1, \dots, \varepsilon_k \alpha_k, u_k)$ is the p -stable normal form of w , then the p -stable normal form of \bar{w} becomes $(\bar{u}_k, -\varepsilon_k \alpha_k, \bar{u}_{k-1}, \dots, -\varepsilon_1 \alpha_1, \bar{u}_0)$.

Example 50 Let $p = a\bar{a}b\bar{a}\bar{a}$ with $b \neq \bar{b}$ and $w = p^4 \bar{b} \bar{a} \bar{a} p^{-1} \bar{a} \bar{a} \bar{b} p^{-2}$. Then the p -stable normal form of w is:

$$(a\bar{a}b\bar{a}\bar{a}, 2, a\bar{a}b\bar{a}\bar{a}\bar{a}\bar{a}, -1, a\bar{a}b\bar{a}\bar{a}\bar{a}\bar{a}, 0, a\bar{a}b\bar{a}\bar{a}).$$

Second case: We assume that \bar{p} is a factor of p^2 . Then we can write $p = rs$ with $\bar{p} = sr$ and $r = \bar{r}$, $s = \bar{s}$. We allow $r = 1$, hence the second case includes the case $p = \bar{p}$. In fact, if $r = 1$, then below we obtain the usual definition

of p -stable normal form, compare e.g. with [3]. The idea is to replace each maximal factor of the form $(rs)^\alpha r$ with $\alpha \geq 2$ by a sequence $rs, \alpha - 2, sr$. In this notation $\alpha - 2$ is representing the factor $(rs)^{\alpha-2} r = p^{\alpha-2} r = r\bar{p}^{\alpha-2} = rp^{2-\alpha}$.

The p -stable normal form (second kind) of $w \in \Gamma^*$ is the shortest sequence (k is minimal)

$$(u_0, \alpha_1, u_1, \dots, \alpha_k, u_k)$$

such that $k \geq 0$, $u_0, u_i \in \Gamma^*$, $\alpha_i \geq 0$ for $1 \leq i \leq k$, and the following conditions are satisfied:

- $w = u_0 p^{\alpha_1} r u_1 \cdots p^{\alpha_k} r u_k$.
- $k = 0$ if and only if $p^2 r$ is not a factor of w .
- If $k \geq 1$, then:

$$\begin{aligned} u_0 &\in \Gamma^* r s \setminus (\Gamma^* p^2 r \Gamma^* \cup \Gamma^* r s r s), \\ u_i &\in (\Gamma^* r s \cap s r \Gamma^*) \setminus (s r s r \Gamma^* \cup \Gamma^* p^2 r \Gamma^* \cup \Gamma^* r s r s) \text{ for } 1 \leq i < k, \\ u_k &\in s r \Gamma^* \setminus (\Gamma^* p^2 r \Gamma^* \cup s r s r \Gamma^*). \end{aligned}$$

Since $\bar{r}\bar{s} = sr$, the p -stable normal form of \bar{w} becomes

$$(\bar{u}_k, \alpha_k, \bar{u}_{k-1}, \dots, \alpha_1, \bar{u}_0).$$

So, for the second kind no negative integers interfere.

Example 51 Let $p = a\bar{a}b$ with $b = \bar{b}$. Then $r = a\bar{a}$ and $s = b$. Let $w = \bar{a}b p^4 a p^3 a$. Then the p -stable normal form of w is:

$$(\bar{a}b a \bar{a} b, 1, b \bar{a} b a a \bar{a} b, 0, b \bar{a} b a).$$

In both cases we can write the p -stable normal form of w as a sequence

$$(u_0, \alpha_1, u_1, \dots, \alpha_k, u_k)$$

where u_i are words and α_i are integers.

It is well-known [28] that for Boolean matrices $A \in \mathbb{B}^{n \times n}$ we have $A^{n!} = A^{n!} A^{n!}$. Hence the matrix $A^{n!}$ is idempotent. For the following we define and fix $c(M_{2n}) = \max\{4, n!\}$. This choice guarantees $h(uv^{c(M_{2n})}w) = h(uv^{2c(M_{2n})}w)$ for all $u, v, w \in \Gamma^*$ and all $h : \Gamma^* \rightarrow M_{2n}$ and, of course, $c(M_{2n}) \geq 3$. The fact $c(M_{2n}) \geq 3$ is used at some point below.

Now, let $w, w' \in \Gamma^*$ be two words whose p -stable normal forms are identical up to the position of the i -th integer. Assume that in the p -stable normal w

at this position there is the integer α_i and that for w' at this position there is α'_i . We know $h(w) = h(w')$ as soon as the following conditions are satisfied: $\alpha_i \cdot \alpha'_i > 0$, $|\alpha_i| \geq c(M_{2n})$, $|\alpha'_i| \geq c(M_{2n})$, and $\alpha_i \equiv \alpha'_i \pmod{c(M_{2n})}$. It is therefore convenient to change the syntax of the p -stable normal form. Each non-zero integer α' is written as $\alpha' = \varepsilon(q + \alpha c(M_{2n}))$ where ε, q, α are uniquely defined by $\varepsilon \in \{+1, -1\}$, $0 \leq q < c(M_{2n})$, and $\alpha \geq 0$. For $\alpha' = 0$ we may choose $\varepsilon = q = \alpha = 0$. The values ε , q , and $c(M_{2n})$ are viewed as constants, if $\alpha = 0$, then it is viewed as a constant, too. Otherwise, if $\alpha \geq 1$, then we view α as a variable ranging over positive integers.

Let u , v , and w be words such that $uv = w$ holds. Write these words in their p -stable normal forms:

$$\begin{aligned} u: & (u_0, \varepsilon_1(q_1 + \alpha_1 c(M_{2n})), u_1, \dots, \varepsilon_k(q_k + \alpha_k c(M_{2n})), u_k), \\ v: & (v_0, \varepsilon'_1(s_1 + \beta_1 c(M_{2n})), v_1, \dots, \varepsilon'_\ell(s_\ell + \beta_\ell c(M_{2n})), v_\ell), \\ w: & (w_0, \varepsilon''_1(t_1 + \gamma_1 c(M_{2n})), w_1, \dots, \varepsilon''_m(t_m + \gamma_m c(M_{2n})), w_m). \end{aligned}$$

Since $uv = w$ there are many identities. For example, for $k, \ell \geq 2$ we have $u_0 = w_0$, $v_\ell = w_m$, $q_1 = t_1$, $\alpha_1 = \gamma_1$, etc. What exactly happens depends only on the p -stable normal form of the product $u_k v_0$. There are several cases, which can be listed easily. We treat only one of them, which is in some sense the most difficult one: We treat the case $p = rs$ with $r = \bar{r}$ and $s = \bar{s}$. It may lead to a large exponent of periodicity. It might be that $u_k = srsr_1$ and $v_0 = r_2srs$ with $r_1 r_2 = r$ (and $r_1 \neq 1 \neq r_2$). Hence we have $u_k v_0 = sp^3$ and $k + \ell = m + 1$. It follows that $\alpha_1 = \gamma_1, \dots, \alpha_{k-1} = \gamma_{k-1}$, $\beta_2 = \gamma_{k+1}, \dots, \beta_\ell = \gamma_m$, and there is only one non-trivial identity:

$$q_k + s_1 + 4 + (\alpha_k + \beta_1)c(M_{2n}) = t_k + \gamma_k c(M_{2n}).$$

Since by assumption $c(M_{2n}) \geq 3$, the case $u_k v_0 = sp^3$ leads to the identity:

$$\gamma_k = \alpha_k + \beta_1 + c \text{ with } c \in \{0, 1, 2\}.$$

Assume now that $\alpha_k \geq 1$ and $\beta_1 \geq 1$. If we replace α_k , β_1 , and γ_k by some $\alpha'_k \geq 1$, $\beta'_1 \geq 1$, and $\gamma'_k \geq 1$ such that we still have $\gamma'_k = \alpha'_k + \beta'_1 + c$, then we obtain new words u' , v' , and w' with the same images under h in M_{2n} and the identity $u'v' = w'$ remains true.

The following step is completely analogous to what has been done in detail in [20, 16, 17, 3]. Using the p -stable normal form we can associate with an equation $L = R$ of denotational length d together with its solution $\sigma : \Omega \rightarrow \Gamma^*$ some linear Diophantine system of d equations in at most $3d$ variables. The variables range over positive natural numbers. The parameters of this system are such that the maximal size of a minimal solution (with respect to the

component wise partial order of \mathbb{N}^d) is in $\mathcal{O}(2^{1.6d})$ with the same approach as in [20]. This tight bound is based in turn on the work of [12]; a more moderate bound $2^{\mathcal{O}(d)}$ (which is enough for our purposes) is easier to obtain, see e.g. [3]. The maximal size of a minimal solution of the linear Diophantine system translates back into a bound on the exponent of periodicity. For this translation we have to multiply the bound using the factor $c(M_{2n})$ and to add $c(M_{2n}) + 1$. Putting everything together we obtain the claim of the proposition since $c(M_{2n}) \in 2^{\mathcal{O}(n \log n)}$. \square

Acknowledgments We thank the anonymous referees and Géraud Sériergues for detailed comments which helped to improve the presentation of the paper. The research has been supported partly by the German Research Foundation *Deutsche Forschungsgemeinschaft, DFG* within the project em GWSS. In addition, Claudio Gutiérrez thanks the Centro de Modelamiento Matemático, FONDAPE Matemáticas Discretas, for financial support.

References

- [1] M. Benois. Parties rationnelles du groupe libre. *C. R. Acad. Sci. Paris, Sér. A*, 269:1188–1190, 1969.
- [2] J. Berstel. *Transductions and context-free languages*. Teubner Studienbücher, Stuttgart, 1979.
- [3] V. Diekert. Makanin’s Algorithm. In M. Lothaire, editor, *Algebraic Combinatorics on Words*, volume 90 of *Encyclopedia of Mathematics and its Applications*, chapter 12, pages 387–442. Cambridge University Press, 2002.
- [4] V. Diekert, C. Gutiérrez, and C. Hagenah. The existential theory of equations with rational constraints in free groups is PSPACE-complete. In A. Ferreira and H. Reichel, editors, *Proc. 18th Annual Symposium on Theoretical Aspects of Computer Science (STACS’01), Dresden (Germany), 2001*, number 2010 in *Lecture Notes in Computer Science*, pages 170–182. Springer-Verlag, 2001.
- [5] V. Diekert and M. Lohrey. Word equations over graph products. In *Proceedings of the 23rd Conference on Foundations of Software Technology and Theoretical Computer Science (FSTTCS 2003), Mumbai (India)*, number 2914 in *Lecture Notes in Computer Science*, pages 156–167. Springer-Verlag, 2003.
- [6] V. Diekert and M. Lohrey. Existential and positive theories of equations in graph products. *Theory Comp. Syst.*, 37:133–156, 2004.
- [7] V. Diekert, Yu. Matiyasevich, and A. Muscholl. Solving word equations modulo partial commutations. *Theoretical Computer Science*, 224:215–235, 1999. Special issue of LFCS’97.
- [8] V. Diekert and A. Muscholl. Solvability of equations in free partially commutative groups is decidable. In F. Orejas, P. G. Spirakis, and J. van Leeuwen, editors, *Proc. 28th International Colloquium on Automata, Languages and Programming (ICALP’01)*, number 2076 in *Lecture Notes in Computer Science*, pages 543–554, Berlin Heidelberg, 2001. Springer-Verlag.
- [9] V. G. Durnev. Undecidability of the positive $\forall\exists^3$ -theory of a free semi-group. *Sibirsky Matematicheskie Jurnal*, 36(5):1067–1080, 1995. In Russian; English translation: *Sib. Math. J.*, 36(5), 917–929, 1995.

- [10] S. Eilenberg. *Automata, Languages, and Machines*, volume A. Academic Press, New York and London, 1974.
- [11] M. R. Garey and D. S. Johnson. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W.H. Freeman and Company, San Francisco, 1979.
- [12] J. von zur Gathen and M. Sieveking. A bound on solutions of linear integer equalities and inequalities. *Proc. Amer. Math. Soc.*, 72(1):155–158, october 1978.
- [13] Y. Gurevich and A. Voronkov. Monadic simultaneous rigid E-unification and related problems. In P. Degano, R. Gorrieri, and A. Marchetti-Spaccamela, editors, *Proc. 24th International Colloquium on Automata, Languages and Programming (ICALP'97)*, Bologna, number 1256 in Lecture Notes in Computer Science, pages 154–165, Berlin Heidelberg, 1997. Springer-Verlag.
- [14] C. Gutiérrez. Satisfiability of word equations with constants is in exponential space. In *Proc. 39th Ann. Symp. on Foundations of Computer Science (FOCS'98)*, Los Alamitos (California), pages 112–119. IEEE Computer Society Press, 1998.
- [15] C. Gutiérrez. Equations in free semigroups with anti-involution and their relation to equations in free groups. In G. H. Gonnet, D. Panario, and A. Viola, editors, *Proceedings Latin American Theoretical INformatics, LATIN'2000*, number 1776 in Lecture Notes in Computer Science, pages 387–396. Springer-Verlag, 2000.
- [16] C. Gutiérrez. Satisfiability of equations in free groups is in PSPACE. In *Proceedings 32nd Annual ACM Symposium on Theory of Computing, STOC'2000*, pages 21–27. ACM Press, 2000.
- [17] C. Hagenah. *Gleichungen mit regulären Randbedingungen über freien Gruppen*. Ph.d.-thesis, Institut für Informatik, Universität Stuttgart, 2000.
- [18] J. E. Hopcroft and J. D. Ullman. *Introduction to Automata Theory, Languages, and Computation*. Addison-Wesley, 1979.
- [19] J. Karhumäki, F. Mignosi, and W. Plandowski. The expressibility of languages and relations by word equations. *Journal of the Association for Computing Machinery*, 47:483–505, 2000.

- [20] A. Kościelski and L. Pacholski. Complexity of Makanin’s algorithm. *Journal of the Association for Computing Machinery*, 43(4):670–684, 1996.
- [21] A. Kościelski and L. Pacholski. Makanin’s algorithm is not primitive recursive. *Theoretical Computer Science*, 191:145–156, 1998.
- [22] D. Kozen. Lower bounds for natural proof systems. In *Proc. of the 18th Ann. Symp. on Foundations of Computer Science, FOCS’77*, pages 254–266, Providence, Rhode Island, 1977. IEEE Computer Society Press.
- [23] M. Lothaire. *Combinatorics on Words*, volume 17 of *Encyclopedia of Mathematics and its Applications*. Addison-Wesley, Reading, MA, 1983. Reprinted by *Cambridge University Press*, 1997.
- [24] G. S. Makanin. The problem of solvability of equations in a free semi-group. *Math. Sbornik*, 103:147–236, 1977. English transl. in *Math. USSR Sbornik* 32 (1977).
- [25] G. S. Makanin. Equations in a free group. *Izv. Akad. Nauk SSR, Ser. Math.* 46:1199–1273, 1983. English transl. in *Math. USSR Izv.* 21 (1983).
- [26] G. S. Makanin. Decidability of the universal and positive theories of a free group. *Izv. Akad. Nauk SSSR, Ser. Mat.* 48:735–749, 1984. In Russian; English translation in: *Math. USSR Izvestija*, 25, 75–88, 1985.
- [27] S. S. Marchenkov. Unsolvability of positive $\forall\exists$ -theory of a free semi-group. *Sibirsky Matematicheskie Jurnal*, 23(1):196–198, 1982. In Russian.
- [28] G. Markowsky. Bounds on the index and period of a binary relation on a finite set. *Semigroup Forum*, 13:253–259, 1977.
- [29] Yu. Matiyasevich. Reduction of trace equations to word equations. Talk given at the “Colloquium on Computability, Complexity, and Logic”, Institut für Informatik, Universität Stuttgart, Germany, Dec. 5–6, 1996.
- [30] Yu. Matiyasevich. Some decision problems for traces. In S. Adian and A. Nerode, editors, *Proceedings of the 4th International Symposium on Logical Foundations of Computer Science (LFCS’97), Yaroslavl, Russia, July 6–12, 1997*, number 1234 in *Lecture Notes in Computer Science*, pages 248–257, Berlin Heidelberg, 1997. Springer-Verlag. Invited lecture.
- [31] Yu. I. Merzlyakov. Positive formulae over free groups. *Algebra i Logika*, 5(4):25–42, 1966. In Russian.

- [32] W. Plandowski. Testing equivalence of morphisms on context-free languages. In J. van Leeuwen, editor, *Algorithms—ESA'94, Second Annual European Symposium*, volume 855 of *Lecture Notes in Computer Science*, pages 460–470, Utrecht, The Netherlands, 1994. Springer.
- [33] W. Plandowski. Satisfiability of word equations with constants is in NEXPTIME. In *Proc. 31st Annual ACM Symposium on Theory of Computing, STOC'99*, pages 721–725. ACM Press, 1999.
- [34] W. Plandowski. Satisfiability of word equations with constants is in PSPACE. In *Proc. 40th Ann. Symp. on Foundations of Computer Science, FOCS'99*, pages 495–500. IEEE Computer Society Press, 1999.
- [35] W. Plandowski. Satisfiability of word equations with constants is in PSPACE. *Journal of the Association for Computing Machinery*, 51:483–496, 2004.
- [36] W. Plandowski and W. Rytter. Application of Lempel-Ziv encodings to the solution of word equations. In K. G. Larsen et al., editors, *Proc. 25th International Colloquium on Automata, Languages and Programming (ICALP'98), Aalborg (Denmark), 1998*, number 1443 in *Lecture Notes in Computer Science*, pages 731–742, Berlin Heidelberg, 1998. Springer-Verlag.
- [37] A. A. Razborov. On systems of equations in a free group. *Izv. Akad. Nauk SSSR, Ser. Mat.* 48:779–832, 1984. In Russian; English translation in: *Math. USSR Izvestija*, 25, 115–162, 1985.
- [38] K. U. Schulz. Makanin's algorithm for word equations — Two improvements and a generalization. In K. U. Schulz, editor, *Word Equations and Related Topics*, number 572 in *Lecture Notes in Computer Science*, pages 85–150, Berlin Heidelberg, 1991. Springer-Verlag.